

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number  
WO 01/92567 A2(51) International Patent Classification: C12Q 1/68.  
A01Q 67/027Birgit [DE/DE]; Heiterwangerstrasse 10, 81373 Munich  
(DE). BECK, Joachim [DE/DE]; Hertenichstrasse 115,  
81477 Munich (DE). HENKEL, Thomas [DE/DE];  
Freienfelsstrasse 20a, 81249 Munich (DE)

(21) International Application Number: PCT/EP01/06165

(22) International Filing Date: 30 May 2001 (30.05.2001)

(74) Agents: LEIDESCHER, Thomas et al.; Zimmermann &  
Partner, Postfach 33 09 20, 80069 Munich (DE).

(25) Filing Language: English

(81) Designated States (national): AU, CA, JP, US.

(26) Publication Language: English

(84) Designated States (regional): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE, TR).(30) Priority Data:  
60/207,400 30 May 2000 (30.05.2000) US(71) Applicant (for all designated States except US): MEDI-  
GENE AG [DE/DE]; Lochhamer Strasse 11, 82152  
Planegg/Martinsried (DE).


Published:

without international search report and to be republished  
upon receipt of that report

(72) Inventors; and

(75) Inventors/Applicants (for US only): BUNK, Daniela  
[DE/DE]; Hofmark 8, 82393 Ilfeldorf (DE). REUNER,For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL TARGET GENES FOR DISEASES OF THE HEART



WO 01/92567 A2

(57) Abstract: The present invention relates to a variety of genes abnormally expressed in heart tissue as well as to fragments of such genes. Assessment of the expression level of these genes may be used for testing the predisposition of mammals and preferably humans for a heart disease or for an acute state of such a disease. Preferred diseases in accordance with the invention are congestive heart failure, dilative cardiomyopathy, hypertrophic cardiomyopathy and ischemic cardiomyopathy. The present invention further relates to methods of identifying compounds capable of normalizing the expression level of the aforementioned genes and of further genes affected by the abnormal expression. The identified compounds may be used for formulating compositions, preferably pharmaceutical compositions for preventing or treating diseases. They may also be used as lead compounds for the development of medicaments having an improved efficiency, a longer half-life, a decreased toxicity etc. and to be employed in the treatment of heart diseases. Included in the invention are also somatic gene therapy methods comprising the introduction of at least one functional copy of any of the above-mentioned genes into a suitable cell. Finally, the invention relates to non-human transgenic animals comprising at least one of the aforementioned genes in their germ line. The transgenic animals of the invention may be used for the development of medicaments for the treatment of heart diseases.

### Novel target genes for diseases of the heart

A variety of documents is cited throughout this specification. The disclosure content of said  
5 documents is herewith incorporated by reference.

The present invention is based on the finding that a variety of genes is abnormally  
expressed in diseased heart tissue. Assessment of the expression level of these genes may  
be used for testing the predisposition of mammals and preferably humans for a heart  
10 disease or for an acute state of such a disease. Diseases that preferably relate to the present  
invention are congestive heart failure, dilative cardiomyopathy, hypertrophic  
cardiomyopathy and ischemic cardiomyopathy. The present invention further relates to  
methods of identifying compounds capable of normalizing the expression level of the  
aforementioned genes and of further genes affected by the abnormal expression. The  
15 identified compounds may be used for formulating compositions, preferably  
pharmaceutical compositions, for preventing or treating diseases. They may also be used as  
lead compounds for the development of medicaments having an improved efficiency, a  
longer half-life, a decreased toxicity etc. and to be employed in the treatment of heart  
diseases. Included in the invention are also somatic gene therapy methods comprising the  
20 introduction of at least one functional copy of any of the above-mentioned genes into a  
suitable cell. Finally, the invention relates to non-human transgenic animals comprising at  
least one of the aforementioned genes in their germ line. The transgenic animals of the  
invention may be used for the development of medicaments for the treatment of heart  
diseases.

25

Referring to studies of the American Heart Association, about 60 million people in the  
USA suffer from Cardiovascular diseases like high blood pressure (50.0 mio), Coronary  
heart disease (12.4 mio), Myocardial infarction (7.3 mio), Angina pectoris (6.4 mio), Stroke  
(4.5 mio), Congenital cardiovascular defects (1.0 mio), and Congestive heart failure (4.7  
30 mio). Hence, it follows that 20 per cent of whole population is affected. The mortality was  
949,619 in 1998 in the USA, which means that about 40 % of all deaths were caused by

Cardiovascular diseases. Since 1900 Cardiovascular diseases are the number one cause of death (1918 was an exception) with one death every 33 seconds on average. At present there is no causal treatment for congestive heart failure available.

- 5 Accordingly, the technical problem underlying the present invention was to provide a new generation of tools useful in the diagnosis, prevention and treatment of heart-related diseases.

The solution to said technical problem is achieved by providing the methods of independent  
10 claims 1, 3, 12, 13, 15, 19, 21, 22, 23, 27, 29, 31, 32, 34, 35, 36, 40 to 44, and 46, the monoclonal antibody according to claim 14, the transgenic non-human mammal according to claim 16, and the use according to independent claim 47. Further advantageous features, aspects and details of the invention are evident from the dependent claims, the description, the examples and the drawings.

15

The invention is based upon the unexpected result that the certain genes coding for the protein sequences given in examples 2 to 11 are deregulated in the comparison of one or more failing heart samples to one or more non-failing heart samples and lead to an upregulation (examples 2, 5, 8, 9, 10) or downregulation (examples 3, 4, 6, 7) of the  
20 described polypeptides measured by their respective mRNAs or cDNAs. The significant changes in gene expression levels suggest a causative role in congestive heart failure.

However, such a causative role for one specific indication of the heart leads to the assumption that a deregulation of such gene(s) might play an important role in other  
25 diseases of the heart as well. Such involvement can easily be tested by methods well known in the art and described e. g. in example 1 of the present application by a comparison of the gene expression levels of such gene between a sample of a healthy mammal and of a mammal having the disease in question. Therefore the subject of this invention does not only relate to dilated cardiomyopathy but also to other diseases of the heart.

30

It is well accepted in the art that upregulation of gene expression of a downregulated target gene by means of a gene therapeutic intervention, compensatory molecules or specific activators, for example of transcription or translation, are potentially very promising therapeutic tools to treat a heart disease that is caused or promoted by the downregulation  
5 of such gene.

On the other hand, downregulation of gene expression and/or protein function of an upregulated target gene by means of specific inhibitors, antisense constructs, ribozymes, antibodies or any other compound (as hereinafter defined) are well accepted tools to treat  
10 a heart disease that is caused or promoted by the upregulation of such gene.

As one gene might be upregulated for one indication of the heart whereas the same gene might be downregulated for another indication of the heart, both upregulation of gene expression as well as downregulation of gene expression and/or protein function might be  
15 useful for the same target gene in different indications.

The same holds true for methods for identifying a subject at risk for a disease of the heart, a method for identifying a compound, a method for identifying one or a plurality of genes as well as methods to make transgenic non-human mammals. In all these various  
20 embodiments of the invention aberrant gene expression in either direction can be used for the given methods.

Thus, the present invention relates to a method for identifying a subject at risk for a disease of the heart, comprising the step of quantitating in the heart tissue of the subject the amount  
25 of at least one RNA encoding an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID  
30 NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8

[66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];

(b) an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a);

5 (c) the amino acid sequence of (a) with at least one conservative amino acid substitution;

(d) an amino acid sequence that is an isoform of the amino acid sequence of any of (a) to (c);

(e) the RNA transcribed from the DNA sequence of SEQ ID NO: 10 [NM\_003970], the DNA sequence of SEQ ID NO: 11 [AW755252], the DNA sequence of SEQ ID NO: 12  
10 [EST clone 52706], the DNA sequence of SEQ ID NO: 13 [EST clone 56461], the DNA sequence of SEQ ID NO: 14 [M14780], the DNA sequence of SEQ ID NO: 15 [61166contig], the DNA sequence of SEQ ID NO: 16 [AF161698], the DNA sequence of SEQ ID NO: 17 [65330contig], the DNA sequence of SEQ ID NO: 18 [66214cds] or the DNA sequence AF129505, or the DNA sequence of SEQ ID NO: 19 [X83703] or a  
15 degenerate variant thereof; and

(f) an amino acid that is encoded by a DNA molecule the complementary strand of which hybridizes in 4xSSC, 0.1% SDS at 65°C to the DNA molecule encoding the amino acid sequence of (a), (c) or (d).

The term "disease of the heart" means, in accordance with the present invention, any  
20 disease that affects the normal function of the heart. This definition includes hereditary as well as acquired diseases such as diseases induced by a pathogen or diseases due to lack of exercise.

Several diseases of the heart are, for example, rheumatic fever/ rheumatic heart disease, hypertensive heart disease, hypertensive heart and renal disease, ischemic heart disease  
25 (coronary heart disease), diseases of pulmonary circulation (which include acute and chronic pulmonary heart disease), arrhythmias, congenital heart disease, angina and congestive heart failure.

The term "quantitating the amount of at least one RNA" is intended to mean the determination of the amount of mRNA in heart tissue as compared to a standard value such  
30 as an internal standard. The (internal) standard would advantageously be the amount of a corresponding RNA produced by a heart tissue not affected by a disease. Said (internal)

standard would also include a mean value obtained from a variety of heart tissues not affected by a disease. A possible way to get samples of heart tissue would be to take a biopsy (catheter) from the ventricular wall. Optionally, a standard would take into account the genetic background of the subject under investigation. Thus, quantitation of said  
5 subject's RNA is effected in comparison to the amount of RNA of one or a variety of samples of the same or a similar genetic background. A variable number of "non-failing" humans (humans that do not show an indication for any heart disease) are compared with a variable number of patients that suffer a distinct heart disease like dilated cardiomyopathy. The determination can be effected by any known technology of analysing the amount of  
10 RNA produced in a sample such as a tissue sample. Techniques based on hybridisation like Northern-Blot, dot-blot, subtractive hybridisation, DNA-Chip analysis or techniques based on reverse transcription coupled to the polymerase chain reaction (RT-PCR) like differential display, suppression subtractive hybridisation (SSH), fluorescence differential display (FDD), serial analysis of gene expression (SAGE) or representational difference  
15 analysis (see e. g. Kozian, D.H., Kirschbaum, B.J.; Comparative gene-expression analysis. (1999) 17:73-77). Generally, it is preferred that the assay is performed as a high throughput assay. This holds also true for the further methods described herein and in accordance with this invention. Samples of RNA may be prepared as described in the appended examples.

20 The term "isoform" means a derivative of a gene resulting from alternative splicing, alternative polyadenylation, alternative promoter usage or RNA editing. Isoforms can be detected by

- (a) *in silico* analysis (e.g. by clustering analysis of any types of expressed sequences or the corresponding proteins, by alignment of expressed sequences with chromosomal DNA,  
25 by interspecies comparisons or by analysis of the coding as well as non-coding sequences like promoters or regulatory RNA processing sites for SNPs or known mutations causing a disease).
- (b) any type of hybridisation techniques (1,2) (e.g. Northern blots, nuclease protection assays, microarrays) starting from RNA.
- 30 (c) PCR-applications as well as hybridisation techniques starting from single strand or double strand cDNA obtained by reverse transcription (3), as described for example in

Higgins, S.J., Hames, D. RNA Processing: A practical approach Oxford University Press (1994), Vol. 1 and 2; Sambrook, Fritsch, Maniatis, Molecular Cloning, a laboratory manual. (1989) Cold Spring Harbor Laboratory Press; Stoss, O. Stoilov, P., Hartmann, A.M., Nayler, O., Stamm, S., The *in vivo* minigene approach to analyse  
5 tissue-specific splicing. Brain Res. Brain Res. Protoc. (1999), 3:383-394.

Primers/probes for RT-PCR or hybridisation techniques are designed in a fashion that at least one of the primers/probes recognizes specifically one isoform. If differences in the molecular weight of isoforms are big enough to separate them with electrophoretical or chromatographical methods, it is also possible to detect multiple isoforms at once by  
10 employing primers/probes that flank the spliced regions. The isoforms are then sequenced and analysed as described in (a).

The term "DNA molecule the complementary strand of which hybridizes in 4xSSC, 0.1% SDS at 65°C to the DNA molecule encoding the amino acid sequence of (a), (c) or (d)"  
15 means that the two DNA molecules hybridize under these experimental conditions to each other. This term does not exclude that the two DNA sequences hybridize at higher stringency conditions such as 2xSSC, 0.1% SDS at 65°C nor does it exclude that lower stringency conditions such as 6xSSC, 0.1% SDS at 60°C allow a hybridization of the two DNA sequences.

20  
Appropriate hybridization conditions for each sequence may be established on well-known parameters such as temperature, composition of the nucleic acid molecules, salt conditions etc.; see, for example, Sambrook et al., "Molecular Cloning, A Laboratory Manual"; CSH Press, Cold Spring Harbor, 1989 or Higgins and Harnes (eds.), "Nucleic acid hybridization,  
25 a practical approach", IRL Press, Oxford 1985, see in particular the chapter "Hybridization Strategy" by Britten & Davidson, 3 to 15.

In accordance with the present invention it has surprisingly been found that a variety of genes is aberrantly expressed in diseases associated with the heart and in particular in  
30 patients suffering from congestive heart failure. By performing the method of the invention which may be *in vivo*, *in vitro* or *in silico*, the diagnosis of a disease of the heart established

by a different methodology may be corroborated. Alternatively, it may be assessed whether a subject that is preferably throughout this specification a human displaying no sign of being affected by a disease of the heart is at risk of developing such a disease. This is possible in cases where the aberrant expression of the gene defined herein above is causative of the disease or is a member of a protein cascade wherein another gene/protein than the one identified herein above is causative for said disease. In this regard, the term "causative" is not limited to mean that the aberrant expression of one gene as identified above or which is a member of said protein cascade is the sole cause for the onset of the disease. Whereas this option is also within the scope of the invention, expression the invention also encompasses embodiments wherein said aberrant is one of a variety of causative events that lead to the onset of the disease.

There is causal correlation between altered cellular function of cardiomyocytes and its protein composition. The latter is regulated by three main mechanisms:

- a. Gene expression
- b. Alternative splicing
- c. Posttranslational modification

In a variation of the method of the invention quantitation of the above recited RNA is used to monitor the progress of a disease of the heart (said variation also applies to the method described herein below). This variation may be employed for assessing the efficacy of a medicament or to determine a time point when administration of a drug is no longer necessary or when the dose of a drug may be reduced and/or when the time interval between administrations of the medicament may be increased. This variation of the method of the invention may successfully be employed in cases where an aberrant expression of any of the aforementioned genes/genes as members of protein cascades is causative of the disease. It is also useful in cases where the aberrant expression of the gene/genes is the direct or indirect result of said disease.

When assessing the risk or the status of the disease, one or more of the RNA levels may be determined. Generally, the assessment of more than 1, such as 2, 3, 4, 5, 6, 7, 8, 9 or 10



different RNAs is expected to enhance the fidelity of the prognosis/diagnosis. However, the gain in fidelity would, as a rule, have to be weighted against the costs generated by such additional tests. Accordingly, it is preferred that one or two different RNA levels are determined for a first assessment. If deemed necessary or appropriate, further RNA levels  
5 may be determined.

In a preferred embodiment of the method of the invention the amount of the said RNA is quantitated using a nucleic acid probe which is a nucleic acid comprising a sequence selected from the group consisting of:

- 10 (a) the DNA sequence of SEQ ID NO: 10 [NM\_003970], the DNA sequence of SEQ ID NO: 11 [AW755252], the DNA sequence of SEQ ID NO: 12 [EST clone 52706], the DNA sequence of SEQ ID NO: 13 [EST clone 56461], the DNA sequence of SEQ ID NO: 14 [M14780], the DNA sequence of SEQ ID NO: 15 [61166contig], the DNA sequence of SEQ ID NO: 16 [AF161698], the DNA sequence of SEQ ID NO: 17 [65330contig], the  
15 DNA sequence of SEQ ID NO: 18 [66214cds] or the DNA sequence AF129505, or the DNA sequence of SEQ ID NO: 19 [X83703] or a degenerate variant thereof (b) a DNA sequence at least 60%, preferably 80%, especially 90%, advantageously 99% identical to the DNA sequence of (a); (c) a nucleic acid sequence that encodes the amino acid sequence  
20 SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid  
25 sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; with at least one conservative amino acid substitution; (d) a nucleic acid sequence that encodes an amino acid sequence that is at least 60%, preferably 80%, especially 90%, advantageously 99% identical to the amino acid sequence of (b); (e) a nucleic acid sequence that encodes the amino acid sequence of (a) or (b) with at least one conservative amino acid substitution;  
30 (f) a nucleic acid sequence that hybridizes in 4xSSC, 0.1% SDS at 65°C to the complementary strand of the DNA molecule encoding the amino acid sequence of (a) or (c); and (g) a fragment of at least 15 nucleotides in length of (a) to (f).

Advantageously, the nucleic acid sequence which is preferably a DNA sequence is detectably labeled. Appropriate labels include radioactive labels, wherein the radioactivity conferring molecules may be, e.g.,  $^{32}\text{P}$ ,  $^{35}\text{S}$  or  $^3\text{H}$ . Appropriate labels further include  
5 fluorescent, phosphorescent or bioluminescent labels or nucleic acid sequences coupled to biotin or streptavidin in order to detect them via anti-biotin or anti-streptavidin antibodies. Whereas any of the above mentioned probes specifically hybridizing to the aforementioned RNAs may be employed, it is preferred that fragments of the full length coding sequence such as oligomers of a length between 15 and 25 nucleotides are used. Examples of such  
10 oligomers are oligomers of 18, 21 or 24 nucleotides. Alternatively, the double strand formed after hybridization can be detected by anti-double strand DNA specific antibodies or aptamers etc.

In this regard, it is understood that the probe of SEQ ID NO: 10 and the mentioned variants thereof are used for quantitating the RNA of SEQ ID NO: 1, but not to any of the other  
15 mentioned RNAs. In the following, appropriate pairs of RNAs and corresponding probes for assessing risks etc. of diseases of the heart are mentioned with the understanding that (i) appropriate variants of the probes as mentioned above may be used and (ii) said probes are specific for the corresponding RNA only but not for any of the other mentioned RNAs. These pairs are: SEQ ID NOs: 2/SEQ ID NO: 11; SEQ ID NO: 3/SEQ ID NO: 13; SEQ ID  
20 NO: 4/SEQ ID NO: 14; SEQ ID NO: 5/SEQ ID NO: 15; SEQ ID NO: 6/SEQ ID NO: 16; SEQ ID NO: 7/SEQ ID NO: 17; SEQ ID NOs: 8/SEQ ID NO: 18; SEQ ID NO: 9/SEQ ID NO: 19.

After hybridization, appropriate washing steps are performed in order to remove unspecific  
25 signals. Appropriate washing conditions include 2 wash steps at 65°C with 2xSSC, 0,1% SDS for 30 min (50 ml) and finally two wash steps with 50 ml of a solution containing 0.1xSSC, 0.1% SDS for 30 min.; see also Sambrook et al., loc. cit., Higgins and Hames, loc. cit. After washing, the label is detected, depending on its nature. For example, a radioactive label may be detected by exposure to an X-ray film or by a phosphorimager.  
30 Alternatively, biotinylated probes can be detected by fluorescence, e.g. by using SAPE (streptavidin-phycoerythrin) with subsequent detection of the signal by a laser scanner.

In addition, the invention relates to a method for identifying a subject at risk for a disease of the heart, comprising the step of quantitating in the heart tissue of the subject the amount of a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid  
5 sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8  
10 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution. Further included are polypeptides  
15 encoded by any of the above recited nucleic acid sequences. This holds also true for any of the other embodiments in which the aforementioned polypeptides are employed.

This embodiment of the invention makes use of the option that detection may not only be at the level of the mRNA but also at the level of the polypeptide translated from the mRNA.  
20 Whereas it is not excluded that the level of mRNA strictly correlates with the level of polypeptide translated from the mRNA, this may not always be the case. Accordingly, it may be assessed whether the mRNA or the protein level, if different, is more appropriate to establish if the heart of a subject is prone to develop a disease of the heart. Factors that contribute to differences in the expression levels of mRNA and protein are well-known in  
25 the art and include differential mRNA-export to the protein-synthesis machinery as well as differences in the translation efficacy of different mRNA species. Other considerations influencing the choice of the detection level (in RNA or protein) include the availability of an appropriate screening tool, instrumentation of the lab, experience of the lab personnel and others.

30

In a preferred embodiment of the method of the invention, the amount of the said

polypeptide is quantitated using an antibody that specifically binds a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid sequence of SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably 80%, especially 90%, advantageously 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution, or an antigen-binding portion of said antibody.

The antibody used in accordance with the invention may be a monoclonal or a polyclonal antibody (see Harlow and Lane, "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, USA, 1988) or a derivative of said antibody which retains or essentially retains its binding specificity. Whereas particularly preferred embodiments of said derivatives are specified further herein below, other preferred derivatives of such antibodies are chimeric antibodies comprising, for example, a mouse or rat variable region and a human constant region. The term "specifically binds" in connection with the antibody used in accordance with the present invention means that the antibody etc. does not or essentially does not cross-react with (poly)peptides of similar structures. Cross-reactivity of a panel of antibodies etc. under investigation may be tested, for example, by assessing binding of said panel of antibodies etc. under conventional conditions (see, e.g., Harlow and Lane, loc. cit.) to the polypeptide of interest as well as to a number of more or less (structurally and/or functionally) closely related polypeptides. Only those antibodies that bind to the polypeptide of interest but do not or do not essentially bind to any of the other (poly)peptides which are preferably expressed by the same tissue as the polypeptide of interest, i.e. heart, are considered specific for the polypeptide of interest and selected for further studies in accordance with the method of the invention.

In a particularly preferred embodiment of the method of the invention, said antibody or antibody binding portion is or is derived from a human antibody or a humanized antibody.

The term "humanized antibody" means, in accordance with the present invention, an antibody of non-human origin, where at least one complementarity determining region (CDR) in the variable regions such as the CDR3 and preferably all 6 CDRs have been replaced by CDRs of an antibody of human origin having a desired specificity. Optionally, the non-human constant region(s) of the antibody has/have been replaced by (a) constant region(s) of a human antibody. Methods for the production of humanized antibodies are described in, e.g., EP-A1 0 239 400 and WO90/07861.

The specifically binding antibody etc. may be detected by using, for example, a labeled secondary antibody specifically recognizing the constant region of the first antibody. However, in a further particularly preferred embodiment of the method of the invention, the antibody, the binding portion or derivative thereof itself is detectably labeled.

Detectable labels include a variety of established labels such as radioactive ( $^{125}\text{I}$ , for example) or fluorescent labels (see, e.g. Harlow and Lane, loc. cit.). Binding may be detected after removing unspecific labels by appropriate washing conditions (see, e.g. Harlow and Lane, loc. cit.).

20

In an additionally preferred embodiment of the method of the invention, said derivative of said antibody is an scFv fragment.

The term "scFv fragment" (single-chain Fv fragment) is well understood in the art and preferred due to its small size and the possibility to recombinantly produce such fragments.

25

In a preferred embodiment of the method of the invention, said RNA is obtained from heart tissue.

A suitable way would be to take a biopsy (catheter) from the ventricular wall. The decision to do this is clearly affected by the severity of the disease and the general constitution of the patient. The cardiologist and the patient have to drive the final decision. In an additionally

30

preferred embodiment of the method of the invention, said polypeptide is quantitated in heart tissue.

5 In another preferred embodiment, the method of the invention further comprises the step of normalizing the amount of RNA against a corresponding RNA from a healthy subject or cells derived from a healthy subject.

The term "healthy subject" means a subject without any indication for heart disease.

The term "normalizing the amount of RNA against a corresponding RNA from a healthy subject or cells derived from a healthy subject" means, in accordance with the present  
10 invention, that levels of mRNA from a comparative number of cells from the heart of said subject under investigation and from the heart of an individual not affected by a disease of the heart are compared. Alternatively, cells from the heart of the subject under investigation may be compared in terms of the indicated mRNA levels with cells derived from the heart of a healthy individual which are kept in cell culture and optionally form a  
15 cell line. Optionally, different sources of cells such as from different individuals and/or different cell lines may be used for the generation of the standard against which the mRNA level of the subject under investigation is compared.

Using the Affymetrix Chip technology, there is also the possibility to use external standards (that are given separately to the hybridisation cocktail) in order to normalize the values of  
20 different oligonucleotide-chips.

In yet another preferred embodiment, the method of the invention further comprises the step of normalizing the amount of polypeptide against a corresponding polypeptide from a healthy subject or cells derived from a healthy subject.

25

The same considerations as developed for the previous embodiment on the mRNA level apply here to the normalization of protein levels.

Additionally, the invention relates to a method for identifying a compound that increases or  
30 decreases the level in heart tissue of a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid sequence of SEQ ID NO: 1 [NP\_003961], the amino

acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution, comprising the steps of: (1) contacting a DNA encoding said polypeptide under conditions that would permit the translation of said polypeptide with a test compound; and (2) detecting an increased or decreased level of the polypeptide relative to the level of translation obtained in the absence of the test compound.

The term "compound" shall mean any biologically active substance that has an effect on heart tissue or a single heart cell, whereas such compound has a positive or negative influence upon such heart tissue or heart cell. Preferred compounds are nucleic acids, preferably coding for a peptide, polypeptide, antisense RNA or a ribozyme or nucleic acids that act independent from their transcription respective their translation as for example as an antisense RNA or ribozyme; natural or synthetic peptides preferably with a relative molecular mass of about 1.000, especially of about 500 peptide analogs polypeptides or compositions of polypeptides, proteins, protein complexes, fusion proteins, preferably antibodies, especially murine, human or humanized antibodies, single chain antibodies, F<sub>ab</sub> fragments or any other antigen binding portion or derivative of an antibody, including modifications of such molecules as for example glycosylation, acetylation, phosphorylation, farnesylation, hydroxylation, methylation or estrification hormones, organic or anorganic molecules or compositions, preferably small molecules with a relative molecular mass of about 1.000, especially of about 500.

The term "under conditions that would permit the translation of said polypeptide" denotes any conditions that allow the in vitro or in vivo translation of the polypeptide of interest. As regards in vitro conditions, translation may be effected in a cell-free system, as described, for example in Stoss, Schwaiger, Cooper and Stamm (1999). J. Biol. Chem. 274: 10951-

10962), using the TNT-coupled reticulocyte lysate system (Promega). With respect to in vivo conditions, physiological conditions such as conditions naturally occurring inside a cell are preferred.

5 Based on the finding that expression of genes encoding the above recited polypeptides is aberrant, the method of the invention allows the convenient identification or isolation of compounds that counteract such aberrant expression such that normal expression levels are restored or essentially restored.

10 The DNA encoding the polypeptide of interest would normally be contained in an expression vector. The expression vectors may particularly be plasmids, cosmids, viruses or bacteriophages used conventionally in genetic engineering that comprise the aforementioned polynucleotide. Preferably, said vector is a gene transfer or targeting vector. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-  
15 associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the polynucleotides into targeted cell population. Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors; see, for example, the techniques described in Sambrook et al., Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989) N.Y. and Ausubel et al., Current Protocols in Molecular  
20 Biology, Green Publishing Associates and Wiley Interscience, N.Y. (1989). Alternatively, the polynucleotides and vectors can be reconstituted into liposomes for delivery to target cells. The vectors containing the polynucleotides can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium phosphate or DEAE-Dextran mediated transfection or electroporation may be used  
25 for eukaryotic cellular hosts; see Sambrook, supra.

Such vectors may comprise further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions. The polynucleotide is operatively linked to expression control sequences allowing expression in eukaryotic cells. Expression of said polynucleotide comprises transcription of the  
30 polynucleotide into a translatable mRNA. Regulatory elements ensuring expression in eukaryotic cells, preferably mammalian cells, are well known to those skilled in the art.



They usually comprise regulatory sequences ensuring initiation of transcription and, optionally, a poly-A signal ensuring termination of transcription and stabilization of the transcript, and/or an intron further enhancing expression of said polynucleotide. Additional regulatory elements may include transcriptional as well as translational enhancers, and/or naturally-associated or heterologous promoter regions. Possible regulatory elements permitting expression in eukaryotic host cells are the AOX1 or GAL1 promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. Furthermore, depending on the expression system used leader sequences capable of directing the polypeptide to a cellular compartment or secreting it into the medium may be added to the coding sequence of the aforementioned polynucleotide and are well known in the art. The leader sequence(s) is (are) assembled in appropriate phase with translation, initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein, or a portion thereof, into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an C- or N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDVI (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3, the Echo<sup>TM</sup> Cloning System (Invitrogen), pSPORT1 (GIBCO BRL) or pRevTet-On/pRevTet-Off or pCI (Promega).

Preferably, the expression control sequences will be eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells. As mentioned above, the vector used in the method of the present invention may also be a gene transfer or targeting vector. Gene therapy, which is based on introducing therapeutic genes into cells by ex-vivo or in-vivo techniques, is one of the most important applications of gene transfer. Suitable vectors and methods for in-vitro or in-vivo gene therapy are described in the literature and

are known to the person skilled in the art; see, e.g., Giordano, *Nature Medicine* 2 (1996), 534-539; Schaper, *Circ. Res.* 79 (1996), 911-919; Anderson, *Science* 256 (1992), 808-813; Isner, *Lancet* 348 (1996), 370-374; Muhlhauser, *Circ. Res.* 77 (1995), 1077-1086; Wang, *Nature Medicine* 2 (1996), 714-716; WO94/29469; WO 97/00957 or Schaper, *Current Opinion in Biotechnology* 7 (1996), 635-640, and references cited therein. The polynucleotides and vectors may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) into the cell. Preferably, said cell is a germ line cell, embryonic cell, or egg cell or derived therefrom, most preferably said cell is a stem cell.

The vector comprising the DNA would be used to transform a suitable eukaryotic host cell. Upon expression of the DNA, which may be constitutive or induced, the test compound would be contacted with the DNA. This can be done by introducing the test compound into the cell. For example, if the test compound is a (poly)peptide, then introduction may be effected by transfection of the corresponding DNA, optionally comprised in a suitable expression vector. If the compound is a small molecule, preferably with a relative molecular weight of up to 1,000, especially up to 500, the introduction into the cell may be effected by direct administration, plus DMSO for hydrophobic compounds, probably liposomal transfer.

In the case that the method of the invention is carried out in vitro, for example, in a cell-free system, then introduction into a cell would not be necessary. Rather, the test compound would be admixed to the in vitro expression system and the effect of said admixture observed.

The effect of the contact of the DNA of interest with the test compound on the protein level may be assessed by any technology that measures changes in the quantitative protein level. Such technologies include Western blots, ELISAs, RIAs and other techniques referred to herein above.

The change in protein level, if any, as a result of the contact of said DNA and said test

compound is compared against a standard. This standard is measured applying the same test system but omits the step of contacting the compound with the DNA. The standard may consist of the expression level of the polypeptide after no compound has been added. Alternatively, the DNA may be contacted with a compound that has previously been demonstrated to have an influence on the expression level.

Compounds tested positive for being capable of enhancing or reducing the amount of polypeptide produced are prime candidates for the direct use as a medicament or as lead compounds for the development of a medicament. Naturally, the toxicity of the compound identified and other well-known factors crucial for the applicability of the compound as a medicament will have to be tested. Methods for developing a suitable active ingredient of a pharmaceutical composition on the basis of the compound identified as a lead compound are described elsewhere in this specification.

Additionally, the invention relates to a method for identifying a compound that specifically binds to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; comprising the steps of (1) providing said polypeptide; and (2) identifying a compound that is capable of binding said polypeptide.

Based on the function of these proteins in DCM development a cell based assay can be developed to identify potential inhibitors or activators. The protein under investigation is expressed in cardiomyocytes (e. g. by infection with recombinant adenovirus). The expression of these proteins lead to characteristic morphological alterations. Reversal or reduction of these morphological alterations can be used in a HTS assay to identify

compounds which act as inhibitors or activators of these proteins. The system can be automated by use of digital image analysis systems.

Another possibility is to identify first proteins which are binding partners of the claimed proteins. This is especially important for structural proteins or adaptor proteins in signal  
5 transduction pathways.

Methods to identify compounds capable of binding are affinity chromatography with immobilised target protein and subsequent elution of bound proteins (e. g. by acid pH), co-immunoprecipitation and as a third method chemical crosslinking with subsequent analysis on SDS-PAGE.

- 10 The influence of compounds on these protein-protein interactions can be monitored by techniques like optical spectroscopy (e. g. fluorescence or surface plasmon resonance), calorimetry (isothermal titration microcalorimetry) and NMR. In the case of optical spectroscopy either the intrinsic protein fluorescence may change (in intensity and/or wavelength of emission maximum) upon complex formation with the binding compound or  
15 the fluorescence of a covalently attached fluorophore may change upon complex formation. The claimed protein or its identified binding partner may be labelled on e. g. cysteine or lysine residues with a fluorophore (for a collection of fluorophores see catalogues of Molecular Probes or Pierce Chemical Company) which changes its optical properties upon binding. These changes in the intrinsic or extrinsic fluorescence may be applied for use in a  
20 HTS assay to identify compounds capable of inhibiting or activating the mentioned protein-protein interaction.

- If the claimed protein exhibits enzymatic activity (e. g. Kinase, Protease, Phosphatase) the inhibition or activation of this activity may be monitored by using labelled (fluorescently, radioactively or immunologically) derivatives of the substrate. This activity assay which is  
25 based on labelled substrates can be used for development of a HTS assay to identify compounds acting as inhibitors or activators.

- Further the invention relates to a monoclonal antibody or derivative thereof that specifically binds to polypeptide having an amino acid sequence selected from the group consisting of  
30 SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID

NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676].

5

Moreover, the invention relates to a method for identifying a compound that increases or decreases the level in heart tissue of an mRNA encoding a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid sequence of SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid  
10 sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a  
15 polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution, the method comprising the steps of (1) contacting a DNA giving rise to said mRNA under conditions that would permit transcription of said  
20 mRNA with a test compound; and (2) detecting an increased or decreased level of the mRNA relative to the level of transcription obtained in the absence of the test compound.

This embodiment of the invention is very similar to the previously discussed one with the exception that here mRNA levels are detected whereas in the previous embodiment protein levels are detected. Methods of assessing RNA levels which also apply to this embodiment  
25 have been described herein above.

Furthermore, the invention relates to a transgenic non-human mammal whose somatic and germ cells comprise at least one gene encoding a functional or disrupted polypeptide selected from the group consisting of: (a) the polypeptide having the amino acid sequence  
30 SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID

NO: 4 [AAAS2025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution, that has been modified, said modification being sufficient to decrease or increase the amount of said functional polypeptide expressed in the heart tissue of said transgenic non-human mammal, wherein said transgenic non-human mammal exhibits a disease of the heart.

A method for the production of a transgenic non-human animal, for example transgenic mouse, comprises introduction of the aforementioned polynucleotide or targeting vector into a germ cell, an embryonic cell, stem cell or an egg or a cell derived therefrom. The non-human animal can be used in accordance with a screening method of the invention described herein. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., *Gene Targeting, A Practical Approach* (1993), Oxford University Press. The DNA of the embryonal membranes of embryos can be analyzed using, e.g., Southern blots with an appropriate probe; see supra. A general method for making transgenic non-human animals is described in the art, see for example WO 94/24274. For making transgenic non-human organisms (which include homologously targeted non-human animals), embryonal stem cells (ES cells) are preferred. Murine ES cells, such as AB-1 line grown on mitotically inactive SNL76/7 cell feeder layers (McMahon and Bradley, *Cell* 62:1073-1085 (1990)) essentially as described (Robertson, E. J. (1987) in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*. E. J. Robertson, ed. (Oxford: IRL Press), p. 71-112) may be used for homologous gene targeting. Other suitable ES lines include, but are not limited to, the E14 line (Hooper et al., *Nature* 326:292-295 (1987)), the D3 line (Doetschman et al., *J. Embryol. Exp. Morph.* 87:27-45 (1985)), the CCE line (Robertson et al., *Nature* 323:445-448 (1986)), the AK-7 line (Zhuang et al., *Cell* 77:875-884 (1994)). The success of generating a mouse line from ES cells bearing a specific targeted mutation depends on the pluripotency of the ES cells (i.

e., their ability, once injected into a host developing embryo, such as a blastocyst or morula, to participate in embryogenesis and contribute to the germ cells of the resulting animal). The blastocysts containing the injected ES cells are allowed to develop in the uteri of pseudopregnant nonhuman females and are born as chimeric mice. The resultant transgenic mice are chimeric for cells having either the recombinase or reporter loci and are backcrossed and screened for the presence of the correctly targeted transgene (s) by PCR or Southern blot analysis on tail biopsy DNA of offspring so as to identify transgenic mice heterozygous for either the recombinase or reporter locus/loci.

The transgenic non-human animals may, for example, be transgenic mice, rats, hamsters, dogs, monkeys, rabbits, pigs, or cows. Preferably, said transgenic non-human animal is a mouse.

In a preferred embodiment of the transgenic non-human mammal of the invention said functional or disrupted gene was introduced into the non-human mammal or an ancestor thereof, at an embryonic stage.

In a further preferred embodiment of the transgenic non-human mammal of the invention the modification is inactivation, suppression or activation of said gene(s) or leads to the reduction or enhancement of the synthesis of the corresponding protein(s).

This embodiment allows for example the study of the interaction of various mutant forms of the aforementioned polypeptides on the onset of the clinical symptoms of a disease related to disorders in the heart. All the applications that have been herein before discussed with regard to a transgenic animal also apply to animals carrying two, three or more transgenes for example encoding different aforementioned nucleic acid molecules. It might be also desirable to inactivate protein expression or function at a certain stage of development and/or life of the transgenic animal. This can be achieved by using, for example, tissue specific, developmental and/or cell regulated and/or inducible promoters which drive the expression of, e.g., an antisense or ribozyme directed against the RNA transcript encoding the corresponding RNA; see also supra. A suitable inducible system is for example tetracycline-regulated gene expression as described, e.g., by Gossen and Bujard (Proc. Natl. Acad. Sci. 89 USA (1992), 5547-5551) and Gossen et al. (Trends

Biotech. 12 (1994), 58-62). Similar, the expression of the mutant protein(s) may be controlled by such regulatory elements.

As mentioned, the invention also relates to a transgenic non-human animal, preferably mammal and cells of such animals which cells contain (preferably stably integrated into their genome) at least one of the aforementioned nucleic acid molecule(s) or part thereof, wherein the transcription and/or expression of the nucleic acid molecule or part thereof leads to reduction of the synthesis of (a) corresponding protein(s). In a preferred embodiment, the reduction is achieved by an anti-sense, sense, ribozyme, co-suppression and/or dominant mutant effect. "Antisense" and "antisense nucleotides" means DNA or RNA constructs which block the expression of the naturally occurring gene product.

Techniques how to achieve this are well known to the person skilled in the art. These include, for example, the expression of antisense-RNA, ribozymes, of molecules which combine antisense and ribozyme functions and/or of molecules which provide for a co-suppression effect; see also supra. When using the antisense approach for reduction of the amount of said proteins in cells, the nucleic acid molecule encoding the antisense-RNA is preferably of homologous origin with respect to the animal species used for transformation. However, it is also possible to use nucleic acid molecules which display a high degree of homology to endogenously occurring nucleic acid molecules encoding such a protein. In this case the homology is preferably higher than 60%, preferably higher than 80%, particularly higher than 90%, more preferably higher than 95% and especially higher than 99%.

In cases where more than one of the aforementioned genes are inactivated, interrelationships of gene products in the onset or progression of the diseases of the heart may be assessed. In this regard, it is also of interest to cross transgenic non-human animals having different transgenes for assessing further interrelationships of gene products in the onset or progression of said disease. Consequently, the offspring of such crosses is also comprised by the scope of the present invention.



In addition, the invention relates to a method for identifying in heart issue a compound that increases or decreases the expression of a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution, said method comprising the steps of: (1) contacting a transgenic non-human mammal as described herein above with a test compound, and (2) detecting an increased/decreased level of expression of said polypeptide relative to the expression in the absence of said test compound.

The test compound which has preferably been tested beforehand for essentially lacking toxicity for the animal can be administered to the animal by any convenient route suitable for administration. These routes include injection, topical and oral administration. Intervals and doses of administration may vary and will be decided upon by the physician/researcher on a case-by-case basis.

Detection, if any, may be effected by a variety of means. For example, if the transgene includes a bioluminescent portion, increase of polypeptide production may be assessed as described, for example, in EP 95 94 1424.4 or in EP 99 12 4640.6. Alternatively, and if the polypeptides are present in the bloodstream, blood of the non-human transgenic animal may be assessed for the changing quantity of the protein. It is preferred in such a case that the gene encoding the polypeptide of interest carries an inducible promoter. Thus, by comparing the situations with and without induction, it can conveniently be determined whether the test compound has indeed an effect on the polypeptide produced or whether the

test compound causes an effect unrelated to the level of polypeptide produced. In certain embodiments of the invention, the non-human transgenic animal will have to be sacrificed in order to assess whether a change in the level of polypeptide expression has occurred. For example, heart tissue may be removed from the sacrificed animal and assessed, using standard technologies, for the expression level of the protein. For example, an antibody specific for the polypeptide may be contacted with the heart tissue and the test developed with a second labeled antibody that is directed to the first antibody. Alternatively, the first antibody itself may be labeled. Heart tissue of a non-human transgenic animal that has been contacted with the test compound would be compared with heart tissue of a non-human transgenic animal that has not been contacted with said test compound.

As mentioned herein above, the transgenic animal may carry more than one of the aforementioned nucleic acid molecules. Accordingly, the effect of a test compound on the expression level of any of these transgenes may be assessed. In addition, a variety of test compounds may be tested, at the same time, for the effect on one or a variety of said transgenes.

A test compound that has proven to be effective in increasing or decreasing the level of the polypeptide of interest and/or in decreasing or increasing the turnover of the polypeptide of interest may be either directly formulated into a medicament (if, for example, its structure is suitable for administration and if it has proven to be non-toxic) or may serve as a lead compound for downstream developments, the results of which may then be formulated into pharmaceutical compositions.

In a preferred embodiment of the method of the invention the test compound prevents or ameliorates a disease of the heart in said transgenic non-human mammal.

In this embodiment, the effect of the test compound may be assessed by observing the disease state of the transgenic animal. Thus, if the animal suffers from a disease of the heart prior to the administration of the test compound and the administration of the test compound results in an amelioration of the disease, then it can be concluded that this test

compound is a prime candidate for the development of a medicament useful also in humans. In addition the compound could also inhibit disease establishment by treatment in advance.

- 5 A further embodiment of the invention is a method for identifying one or a plurality of isogenes of a gene coding for a polypeptide selected from the group consisting of: the amino acid sequence of SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; the method comprising the steps of
- 10 (1) providing nucleic acid coding for said polypeptide or a part thereof; and
- 15 (2) identifying a second nucleic acid that (i) has a homology of 60%, preferably 80%, especially 90%, advantageously 99% or (ii) hybridizes in 4xSSC, 0.1 SDS at 45°C to the nucleic acid molecule encoding the amino acid sequence of (a), (c) or (d).

The term isogenes shall mean genes that are thought to be created by gene duplication.

20 They can be identified by comparing the homology of the DNA-, RNA-, or protein-sequence of interest with other DNA, RNA or protein-sequences of the same species from different databases. There might be strong differences in the degree of homology between isogenes of the same species. This may be dependent on the time-point, when the gene duplication event took place in evolution and the degree of conservation during evolution.

25 Isogenes can be identified and cloned by RT-PCR as has been demonstrated by Screaton *et al.* (1995) EMBO J. 14:4336-4349 or Huang *et al.* (1998) Gene 211: 49-55. Isogenes can also be identified and cloned by colony hybridisation or plaque hybridization as described in Sambrook, Fritsch, Maniatis (1989), Molecular Cloning. Cold Spring Harbor Laboratory Press. In a first step, either a genomic or a cDNA library in bacteria or phages is generated.

30 In order to identify isogenes, colony hybridisation or plaque hybridization is slightly

modified in a way that cross-hybridizations are detectable under conditions of lower stringency. This can be achieved by lowering the calculated temperature for hybridisation and washing and/or by lowering the salt concentration of the washing solutions (Sambrook, Fritsch, Maniatis (1989) Cold Spring Harbor Laboratory Press). For example, a low-stringency washing condition may include 2 wash steps at a temperature between 45°C and 65°C with 4xSSC, 0.1% SDS for 30 min (50 ml) and finally two wash steps with 50 ml of a solution containing 2xSSC, 0.1% SDS for 30 min. After detection, signal intensity of colonies containing an isogene is dependent on the homology of a gene and its isogene(s).

- Furthermore, the invention relates to a method for identifying one or a plurality of genes whose expression in heart tissue is modulated by inhibiting, decreasing or increasing the expression of a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution, or of an mRNA encoding said polypeptide, said modulation being indicative of a disease of the heart, said method comprising the steps of: (1) contacting a plurality of heart tissue cells with a compound that inhibits, decreases or increases the expression of said polypeptide under conditions that permit the expression of said polypeptide in the absence of a test compound, and (2) comparing a gene expression profile of said heart cell in the presence and in the absence of said compound.

The term "gene expression profile" shall mean all expressed genes of a cell or a tissue.

- Such profile can be assessed using the methods well known in the art, for example isolation of total RNA, isolation of poly(A) RNA from total RNA, suppression subtractive

hybridization, differential display, preparation of cDNA libraries or quantitative dot blot analysis, as for example described in Example 1 of this application.

This embodiment of the method of the invention is particularly suitable for identifying further genes the expression level of which is directly affected by the aberrant expression of  
5 any of the aforementioned genes. In other words, this embodiment of the method of the invention allows the identification of genes involved in the same protein cascade as the aberrantly expressed gene. Typically, the method of the invention will be a method performed in cell culture.

10 The method of the invention allows for the design of further medicaments that use other targets than the aberrantly expressed gene. For example, if a potential target downstream of the aberrantly expressed gene is indeed targeted by a medicament, the negative effect of the aberrantly expressed gene may be efficiently counterbalanced. Compounds modulating other genes in the cascade may have to be refined or further developed prior to  
15 administration as a medicament as described elsewhere in this specification.

Additionally, the invention relates to a method for identifying one or a plurality of genes whose expression in heart tissue is modulated by the inhibition, decreasing or increasing of the expression of a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ  
20 ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID  
25 NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution, or of an mRNA encoding said polypeptide, said modulation being indicative of a disease of the heart, said method  
30 comprising the steps of: (1) providing expression profiles of (i) a plurality of heart tissue cells from or derived from a heart of a subject suffering from a disease of the heart; and (ii)

a plurality of heart tissue cells from or derived from a subject not suffering from a disease of the heart; and (2) comparing the expression profiles (i) and (ii).

In variation to the method described herein above, this embodiment of the method of the invention compares the expression profiles of cells from a healthy subject and a subject suffering from a heart disease. In this regard, the term "cells derived from a heart" includes cells that are held in cell culture or even cell lines that autonomously grow in cell culture and that were originally derived from heart tissue. By comparing the two expression profiles, differences in expression levels of genes involved in the disease of the heart may be identified. As with the preceding embodiment, these genes may be part of a cascade involving the aberrantly expressed gene. Examples of such cascades are signaling cascades. Once genes are identified that are expressed at a different level in a diseased heart, they may be tested up-regulation or down-regulation by bringing them into contact with suitable test compounds. Again, these test compounds may then, with or without further development, be formulated into pharmaceutical compositions.

In a preferred embodiment, the method of the invention further comprises the steps of (3) determining at least one gene that is expressed at a lower or higher level in the presence of said compound; and (4) identifying a further compound that is capable of raising or lowering the expression level of said at least one gene.

This preferred embodiment of the invention requires that one of the genes the expression of which may directly or indirectly be lowered or increased by the expression of the aberrant gene is identified. Then, a further panel of test compounds may be tested for the capacity to increase or decrease the expression of said further gene. Compounds that are successfully tested would be prime candidates for the development of medicaments for the prevention or treatment of a disease of the heart.

In another preferred embodiment, the method of the invention further comprises the steps of (3) determining at least one gene that is expressed at a lower or higher level in said heart tissue cells from or derived from a heart of a subject suffering from a disease of the heart;

and (4) identifying a further compound that is capable of raising or lowering the expression level of said at least one gene.

In variation of the previously discussed embodiment, this embodiment requires that at least one gene is identified by comparing the expression profiles of tissue or cells derived from a healthy subject and from a subject suffering from a disease of the heart. Subsequently, at least one compound is identified that is capable of increasing or decreasing the expression of said gene.

In yet another preferred embodiment, the method of the invention further comprises the steps of (3) determining at least one gene that is expressed at a higher or lower level in the presence of said compound; and (4) identifying a further compound that is capable of reducing or raising the expression level of said at least one gene.

In this and the following embodiment, the situation is covered that another gene in the cascade that also includes the aberrantly expressed gene has a higher or lower expression level that needs to be lowered or raised in order to effectively treat the disease of the heart. Again, once such a gene is identified, a compound is tested for its capacity to lower expression of said gene.

In still another preferred embodiment, the method of the invention further comprises the steps of (3) determining at least one gene that is expressed at a higher or lower level in said heart tissue cells from or derived from a heart of a subject suffering from a disease of the heart; and (4) identifying a further compound that is capable of reducing or enhancing the expression level of said at least one gene.

Additionally, the invention relates to a method for identifying proteins or a plurality of proteins whose activity is modulated by a polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO: 1 [NP 003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid

sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAAS8676]; the method comprising the steps of

5 (1) providing said polypeptide and (2) identifying a further protein that is capable of interacting with said polypeptide.

One possible method to identify protein-protein interactions is the Yeast two-hybrid screen described by Golemis & Khazak (1997), Methods Mol Biol. 63:197-218. Other well

10 established methods in order to identify protein-protein interactions are co-immunoprecipitations or *in vitro* protein interaction assays like GST-pulldown assays (such as described in Stoss, Schwaiger, Cooper and Stamm (1999). J. Biol. Chem. 274: 10951-10962).

15 In a further preferred embodiment of the method of the invention said compound is a small molecule or a peptide derived from an at least partially randomized peptide library.

Additionally, the invention relates to a method of refining a compound identified by the method as described herein above comprising the steps of (1) identification of the binding

20 sites of the compound and the DNA or mRNA molecule by site-directed mutagenesis or chimeric protein studies; (2) identification of the binding-site of said polypeptide and the compound by site-directed mutagenesis of the corresponding DNA or by chimeric protein studies, (3) molecular modeling of both the binding site of the compound and the binding site of the DNA or mRNA molecule; and (4) modification of the compound to improve its

25 binding specificity for the DNA or mRNA.

All techniques employed in the various steps of the method of the invention are conventional or can be derived by the person skilled in the art from conventional techniques without further ado. Thus, biological assays based on the herein identified nature of the

30 polypeptides may be employed to assess the specificity or potency of the drugs wherein the increase of one or more activities of the polypeptides may be used to monitor said



specificity or potency. Steps (1) and (2) can be carried out according to conventional protocols. A protocol for site directed mutagenesis is described in Ling MM, Robinson BH. (1997) Anal. Biochem. 254: 157-178. The use of homology modelling in conjunction with site-directed mutagenesis for analysis of structure-function relationships is reviewed in Szklarz and Halpert (1997) Life Sci. 61:2507-2520. Chimeric proteins are generated by ligation of the corresponding DNA fragments via a unique restriction site using the conventional cloning techniques described in Sambrook, Fritsch, Maniatis. Molecular Cloning, a laboratory manual. (1989) Cold Spring Harbor Laboratory Press. A fusion of two DNA fragments that results in a chimeric DNA fragment encoding a chimeric protein can also be generated using the gateway-system (Life technologies), a system that is based on DNA fusion by recombination. A prominent example of molecular modelling is the structure-based design of compounds binding to HIV reverse transcriptase that is reviewed in Mao, Sudbeck, Venkatachalam and Uckun (2000). Biochem. Pharmacol. 60: 1251-1265.

For example, identification of the binding site of said drug by site-directed mutagenesis and chimerical protein studies can be achieved by modifications in the (poly)peptide primary sequence that affect the drug affinity; this usually allows to precisely map the binding pocket for the drug.

As regards step (2), the following protocols may be envisaged: Once the effector site for drugs has been mapped, the precise residues interacting with different parts of the drug can be identified by combination of the information obtained from mutagenesis studies (step (1)) and computer simulations of the structure of the binding site provided that the precise three-dimensional structure of the drug is known (if not, it can be predicted by computational simulation). If said drug is itself a peptide, it can be also mutated to determine which residues interact with other residues in the polypeptide of interest.

Finally, in step (3) the drug can be modified to improve its binding affinity or its potency and specificity. If, for instance, there are electrostatic interactions between a particular residue of the polypeptide of interest and some region of the drug molecule, the overall charge in that region can be modified to increase that particular interaction.

30

Identification of binding sites may be assisted by computer programs. Thus, appropriate

computer programs can be used for the identification of interactive sites of a putative inhibitor and the polypeptide by computer assisted searches for complementary structural motifs (Fassina, Immunomethods 5 (1994), 114-120). Further appropriate computer systems for the computer aided design of protein and peptides are described in the prior art, for example, in Berry, Biochem. Soc. Trans. 22 (1994), 1033-1036; Wodak, Ann. N. Y. Acad. Sci. 501 (1987), 1-13; Pabo, Biochemistry 25 (1986), 5987-5991. Modifications of the drug can be produced, for example, by peptidomimetics and other inhibitors can also be identified by the synthesis of peptidomimetic combinatorial libraries through successive chemical modification and testing the resulting compounds. Methods for the generation and use of peptidomimetic combinatorial libraries are described in the prior art, for example in Ostresh, Methods in Enzymology 267 (1996), 220-234 and Dorner, Bioorg. Med. Chem. 4 (1996), 709-715. Furthermore, the three-dimensional and/or crystallographic structure of activators of the expression of the polypeptide of the invention can be used for the design of peptidomimetic activators, e.g., in combination with the (poly)peptide of the invention (Rose, Biochemistry 35 (1996), 12933-12944; Rutenber, Bioorg. Med. Chem. 4 (1996), 1545-1558).

In accordance with the above, in a preferred embodiment of the method of the invention said compound is further refined by peptidomimetics.

20

The invention furthermore relates to a method of modifying a compound identified or refined by the method as described herein above as a lead compound to achieve (1) modified site of action, spectrum of activity, organ specificity, and/or (2) improved potency, and/or (3) decreased toxicity (improved therapeutic index), and/or (4) decreased side effects, and/or (5) modified onset of therapeutic action, duration of effect, and/or (6) modified pharmacokinetic parameters (resorption, distribution, metabolism and excretion), and/or (7) modified physico-chemical parameters (solubility, hygroscopicity, color, taste, odor, stability, state), and/or (8) improved general specificity, organ/tissue specificity, and/or (9) optimized application form and route by (i) esterification of carboxyl groups, or (ii) esterification of hydroxyl groups with carbon acids, or (iii) esterification of hydroxyl groups to, e.g. phosphates, pyrophosphates or sulfates or hemi succinates, or (iv) formation

30

of pharmaceutically acceptable salts, or (v) formation of pharmaceutically acceptable complexes, or (vi) synthesis of pharmacologically active polymers, or (vii) introduction of hydrophylic moieties, or (viii) introduction/exchange of substituents on aromates or side chains, change of substituent pattern, or (ix) modification by introduction of isosteric or bioisosteric moieties, or (x) synthesis of homologous compounds, or (xi) introduction of branched side chains, or (xii) conversion of alkyl substituents to cyclic analogues, or (xiii) derivatisation of hydroxyl group to ketals, acetals, or (xiv) N-acetylation to amides, phenylcarbamates, or (xv) synthesis of Mannich bases, imines, or (xvi) transformation of ketones or aldehydes to Schiff's bases, oximes, acetals, ketals, enolesters, oxazolidines, thiozolidines or combinations thereof.

The various steps recited above are generally known in the art. They include or rely on quantitative structure-action relationship (QSAR) analyses (Kubinyi, "Hansch-Analysis and Related Approaches". VCH Verlag, Weinheim, 1992), combinatorial biochemistry, classical chemistry and others (see, for example, Holzgrabe and Bechtold, Deutsche Apotheker Zeitung 140(8), 813-823, 2000).

The invention additionally relates to a method for inducing a disease of the heart in a non-human mammal, comprising the step of contacting the heart tissue of said mammal with a compound that inhibits, decreases or increases the expression of a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution.

This embodiment of the invention is particularly useful for mimicking factors/developments leading to the onset of the disease. The fact, that differences in the expression of a protein contributes to heart failure has been shown for phospholamban, for example. Mice over-expressing phospholamban develop heart failure. This effect is thought to be due to the inhibition of Serca. (Minamisawa et al. (1999) Cell, 99:313-322).

In a preferred embodiment of the method of the invention said compound that decreases or increases is a small molecule, an antibody or an aptamer that specifically binds said polypeptide.

The terms "small molecule" as well as "antibody" have been described herein above and bear the same meaning in connection with this embodiment.

The invention moreover relates to a method of producing a pharmaceutical composition comprising formulating the compound identified, refined or modified by the method as described herein above, optionally with a pharmaceutically active carrier and/or diluent.

The pharmaceutical composition of the present invention may further comprise a pharmaceutically acceptable carrier and/or diluent. Examples of suitable pharmaceutical carriers are well known in the art and include phosphate buffered saline solutions, water, emulsions, such as oil/water emulsions, various types of wetting agents, sterile solutions etc. Compositions comprising such carriers can be formulated by well known conventional methods. These pharmaceutical compositions can be administered to the subject at a suitable dose. Administration of the suitable compositions may be effected by different ways, e.g., by intravenous, intraperitoneal, subcutaneous, intramuscular, topical, intradermal, intranasal or intrabronchial administration. The dosage regimen will be determined by the attending physician and clinical factors. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. A typical dose can be, for example, in the range of 0.001 to 1000 µg (or of nucleic acid for

expression or for inhibition of expression in this range); however, doses below or above this exemplary range are envisioned, especially considering the aforementioned factors. Generally, the regimen as a regular administration of the pharmaceutical composition should be in the range of 1  $\mu$ g to 10 mg units per day. If the regimen is a continuous infusion, it should also be in the range of 1  $\mu$ g to 10 mg units per kilogram of body weight per minute, respectively. Progress can be monitored by periodic assessment. Dosages will vary but a preferred dosage for intravenous administration of DNA is from approximately 10<sup>6</sup> to 10<sup>12</sup> copies of the DNA molecule. The compositions of the invention may be administered locally or systemically. Administration will generally be parenterally, e.g., intravenously; DNA may also be administered directly to the target site, e.g., by biolistic delivery to an internal or external target site or by catheter to a site in an artery. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like. Furthermore, the pharmaceutical composition of the invention may comprise further agents such as interleukins or interferons depending on the intended use of the pharmaceutical composition.

The invention also relates to a method for preventing or treating a disease of the heart in a subject in need of such treatment, comprising the step of increasing or decreasing the level of a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid

sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution, in the heart tissue of the subject.

Further, the invention relates to a method of preventing or treating a disease of the heart in a subject in need of such treatment comprising the step of increasing or decreasing the level of mRNA encoding a polypeptide selected from the group consisting of: (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution, in the heart tissue of the subject.

The invention in a preferred embodiment relates to a method wherein such increase/decrease is effected by administering the pharmaceutical composition obtained by the method as described herein above.

In a further preferred embodiment of the method of the invention such an increase/decrease is effected by introducing the DNA sequence recited herein above into the germ line or into somatic cells of a subject in need thereof.

Technologies for effecting such an introduction have been described herein above.

In a most preferred embodiment of the method of the invention, the disease of the heart to be treated is congestive heart failure, dilative cardiomyopathy, hypertrophic

cardiomyopathy, ischemic cardiomyopathy, specific heart muscle disease, rhythm and conduction disorders, syncope and sudden death, coronary heart disease, systemic arterial hypertension, pulmonary hypertension and pulmonary heart disease, valvular heart disease, congenital heart disease, pericardial disease or endocarditis.

5

In addition, the invention relates to a method for identifying subjects at risk for heart diseases, especially congestive heart failure comprising the step of detecting an increased level of MYOM2, the LIM domain, the muscle isoform of creatine kinase, YAP65, APOBEC-2, SMPX or C-193 (CARP) in the heart tissue of a subject.

10

The invention additionally relates to a method for preventing or treating heart diseases, especially congestive heart failure in a subject, said method comprising the step of contacting the heart tissue of said subject with a compound that decreases or increases the expression of MYOM2, the LIM domain, the muscle isoform of creatine kinase, YAP65, APOBEC-2, SMPX or C-193 (CARP).

15

In addition the invention relates to a method for identifying subjects at risk for heart diseases, especially congestive heart failure comprising the step of detecting decreased creatine kinase activity in the tissue of a subject, especially in a muscle tissue or from blood or serum. One possible method to detect the activity of creatine kinase would be a conventional kinetic UV-test as described by the International Federation of Clinical Chemistry and Laboratory Medicine (IFCC), 1991.

20

Moreover the invention relates to a method for identifying a subject at risk for heart diseases, especially congestive heart failure, said method comprising detecting increased levels of creatine phosphate in a subject, especially in the blood or serum of a subject.

25

The invention as well relates to a method for preventing or treating heart diseases, especially congestive heart failure in a subject, said method comprising the step of increasing the transfer of phosphoryl groups from creatine phosphate to ADP in the tissue of a subject, especially in a muscle tissue.

30

In a preferred embodiment of the method of the invention the activity of creatine kinase is increased in said tissue.

- 5 The invention additionally relates to a method for identifying a compound for preventing or treating heart diseases, especially congestive heart failure, said method comprising the steps of (a) contacting creatine kinase with a substrate for creatine kinase and a test compound, and (b) determining whether the transfer of phosphoryl groups from the substrate is increased in the presence of the test compound.

10

The figures show:

Fig. 1 a shows the cDNA sequence of clone 40399 (corresponds to SEQ ID NO: 20).

- 15 Fig. 1 b shows the sequence of the EST clone NM\_003970. Start and stop codons are marked by bold letters, the sequence of 40399 is marked in italic letters (corresponds to SEQ ID NO: 10).

Fig. 1 c shows the putative amino acid sequence M-PROTEIN (MYOMESIN) 2 (MYOM2) (corresponds to SEQ ID NO: 1).

20

Fig. 1 d shows a schematic alignment of the cDNA fragment 40399 identified in SSH with its homologous Genbank entree and the open reading frame of 1465 amino acids (aa). Not to scale. Homology scores were determined using blast2 algorithm of NCBI:

25

40399-NM\_003970: Expect = 2e-88, Identities = 187/194 (96%), Positives = 187/194 (96%).

Fig. 1 e: Two filters were hybridized sequentially with [ $\alpha$ -33P]UTP labeled T3 transcripts from cDNA libraries prepared from mRNA of five control and

30



four DCM heart tissues as indicated. Experiments were normalized by adjusting the overall signal intensity of each hybridization to 100%, relative expression levels are given. A mean value and standard deviation was calculated from all NF samples and DCM sample 15 and 13, respectively. Asterisks mark samples used for SSH.

Fig. 2 a shows the cDNA sequence of clone 41441 (corresponds to SEQ ID NO: 2).

Fig. 2 b shows the sequence of the EST clone AW755252 (corresponds to SEQ ID NO: 11). Start and stop codons are marked in bold letters, the sequence of 41441 is given in italic letters.

Fig. 2 c shows the amino acid sequence 41441pep (corresponds to SEQ ID NO: 21). The first methionine of the open reading frame is marked in bold letters. Amino acids 11-62 of 41441pep encode a cysteine-rich LIM domain (PS00478, PS50023), which is composed of 2 special zinc fingers that are joined by a 2-amino acid spacer (consensus: CX2CX15-21[FYWH]HX2[CH]X2CX2CX3[LIVMF]XnCX2H as underlined). According to this analyses, we expect the start codon to be further upstream of the first methionine in frame 1 assuming that a sequencing error exists in the 5' region of AW755252.

Fig. 2 d shows a schematic alignment of the cDNA fragment 41441 identified in SSH with its homologous Genbank entree and the predicted open reading frame. Not to scale. Homology scores were determined using blast2 algorithm of NCBI:

41441-AW755252: Expect = 0.0, Identities = 369/385 (95%), Positives = 369/385 (95%), Gaps = 2/385 (0%)

- Fig. 2 e: Two filters were hybridized sequentially with [ $\alpha$ -33P]UTP labeled T3 transcripts from cDNA libraries prepared from mRNA of five control and four DCM heart tissues as indicated. Experiments were normalized by adjusting the overall signal intensity of each hybridization to 100%, relative expression levels are given. Mean values and standard deviations were calculated from all NF and DCM samples, respectively. Asterisks mark samples used for SSH.
- Fig. 3 a shows the cDNA sequence of clone 52706 (corresponds to SEQ ID NO: 12).
- Fig. 3 b: Two filters were hybridized sequentially with [ $\alpha$ -33P]UTP labeled T3 transcripts from cDNA libraries prepared from mRNA of five control, and five DCM heart tissues as indicated. Experiments were normalized by adjusting the overall signal intensity of each hybridization to 100%, relative expression levels are given.
- Fig. 4 a shows the cDNA sequence of clone 56461 (corresponds to SEQ ID NO: 13).
- Fig. 4 b shows the sequence of the EST clone AF077035 (corresponds to SEQ ID NO: 22). Start and stop codons are marked in bold letters, the sequence of 56461 is marked in italic letters.
- Fig. 4 c shows the putative amino acid sequence AAD27768 (corresponds to SEQ ID NO: 3). The first methionine of the open reading frame is marked in bold letters. Amino acids 27-79 of 56461 are highly homologous to the rRNA binding motif of 30S ribosomal protein S17 and 40S ribosomal protein S11 (PD001295). A cleavage site for mitochondrial presequences may be predicted for amino acids 57-61 KRK|TY (R2-motif).

Fig. 4 d shows a schematic alignment of the cDNA fragment 56461 identified in SSH with its homologous Genbank entree and the open reading frame of 130 amino acids (aa). Not to scale. Homology scores were determined using blast2 algorithm of NCBI:

5 56461-AF077035: Expect = 0.0, Identities = 498/502 (99%), Positives = 498/502 (99%), Gaps = 2/502 (0%).

Fig. 4 e: Two filters were hybridized sequentially with [ $\alpha$ -33P]UTP labeled T3 transcripts from cDNA libraries prepared from mRNA of five control and  
10 five DCM heart tissues as indicated. Experiments were normalized by adjusting the overall signal intensity of each hybridization to 100%, relative expression levels are given. A mean value and standard deviation was calculated from all NF samples and DCM15 and DCM13, respectively.

15 Fig. 5 a shows the cDNA sequence of clone 61105 (corresponds to SEQ ID NO: 23).

Fig. 5 b shows the sequence of the EST clone M14780 (corresponds to SEQ ID NO: 14). Start and stop codons are marked by bold letters, the sequence of 61105 is marked in italic letters.

20

Fig. 5 c shows the putative amino acid sequence AAA52025 (corresponds to SEQ ID NO: 4).

Fig. 5 d shows a schematic alignment of the cDNA fragment 61105 identified in SSH with its homologous Genbank entree and open reading frame of 381 amino  
25 acids (aa). Not to scale. Homology scores were determined using blast2 algorithm of NCBI:

61105-M14780: Expect = 0.0, Identities = 375/379 (98%), Positives = 375/379 (98%), Gaps = 1/379 (0%).

30

- Fig. 5 c: Two filters were hybridized sequentially with [ $\alpha$ -<sup>32</sup>P]UTP labeled T3 transcripts from cDNA libraries prepared from mRNA of five control heart tissues and five DCM heart tissues as indicated. Experiments were normalized by adjusting the overall signal intensity of each hybridization to 100%, relative expression levels are given. Mean values and standard deviations were calculated from relative expression levels.
- Fig. 6 a shows the cDNA sequence of clone 61166 (corresponds to SEQ ID NO: 24).
- Fig. 6 b shows the sequence 61166contig assembled from overlapping EST sequences, which are available from public databases (corresponds to SEQ ID NO: 15). Start and stop codons are marked by bold letters, the sequence of 61166 is marked in italic letters.
- Fig. 6 c shows the amino acid sequence of 61166pep (corresponds to SEQ ID NO: 5) Amino acids 40-46 of 61166pep encode a nuclear localization signal pattern 7 (PX1-3[KR][KR][KR], underlined) not present in human YAP65 (NP\_006097). Therefore this protein is expected to be located in the nucleus.
- Fig. 6 d shows a schematic alignment of the cDNA fragment 61166 identified in SSH with its overlapping contig of assembled EST sequences according to LabOnWeb (Compugen) analysis, accession numbers of homologous Genbank entrees and the longest open reading frame of 398 amino acids (aa). Not to scale. Homology scores were determined using blast2 algorithm of NCBI:
- Contig-61166: Expect = 0.0, Identities = 401/403 (99%), Positives = 401/403 (99%), Gaps = 1/403 (0%)
- Contig-AL050107: Expect = 0.0, Identities = 3058/3098 (98%), Positives = 3058/3098 (98%)

Contig-A1927050: Expect = 0.0. Identities = 532/532 (100%), Positives = 532/532 (100%)

Contig-A1745235: Expect = 0.0. Identities = 557/573 (97%), Positives = 557/573 (97%), Gaps = 1/573 (0%)

5

Fig. 6 e: Two filters were hybridized sequentially with [ $\alpha$ -33P]UTP labeled T3 transcripts from cDNA libraries prepared from mRNA of five control heart tissues and five DCM heart tissues as indicated. Experiments were normalized by adjusting the overall signal intensity of each hybridization to 100%, relative expression levels are given. Mean values and standard deviations are given on the right side. Asterisks mark samples used for SSH.

10

Fig. 7 a shows the cDNA sequence of clone 61244 (corresponds to SEQ ID NO: 25)

15

Fig. 7 b shows the sequence of the EST clone AF161698 (corresponds to SEQ ID NO: 16). Start and stop codons are marked by bold letters, the sequence of 61244 is marked in italic letters.

20

Fig. 7 c shows the putative amino acid sequence AAD45360 (corresponds to SEQ ID NO: 6).

25

Fig. 7 d shows a schematic alignment of the cDNA fragment 61244 identified in SSH with its homologous Genbank entree and open reading frame of 224 amino acids (aa). Not to scale. Homology scores were determined using blast2 algorithm of NCBI:

61244-AF161698: Expect = 3e-86, Identities = 168/168 (100%), Positives = 168/168 (100%).

30

Fig. 7 e: Two filters were hybridized sequentially with [ $\alpha$ -33P]UTP labeled T3 transcripts from cDNA libraries prepared from mRNA of five control heart

tissues and five DCM heart tissues as indicated. Experiments were normalized by adjusting the overall signal intensity of each hybridization to 100%, relative expression levels are given. Mean values and standard deviations were calculated from relative expression levels. Asterisks mark samples used for SSH.

Fig. 8 a shows the cDNA sequence of clone 65330 (corresponds to SEQ ID NO: 26).

Fig. 8 b shows the contig of assembled EST sequences (corresponds to SEQ ID NO: 17). Start and stop codons are marked by bold letters, the sequence of 65330 is marked in italic letters.

Fig. 8 c shows the putative amino acid sequence of clone 65330 (corresponds to SEQ ID NO: 7).

Fig. 8 d shows a schematic alignment of the cDNA fragment 65330 identified in SSH with its overlapping contig of assembled EST sequences according to LabOnWeb (Compugen) analysis, accession numbers of homologous Genbank entree and the longest open reading frame of 264 amino acids (aa). Not to scale. Homology scores were determined using blast2 algorithm of NCBI:

Contig-65330: Expect = 0.0, Identities = 334/334 (100%), Positives = 334/334 (100%)

Contig-AF249873: Expect = 0.0, Identities = 1020/1028 (99%), Positives = 1020/1028 (99%).

Fig. 8 e: Two filters were hybridized sequentially with [ $\alpha$ -33P]UTP labeled T3 transcripts from cDNA libraries prepared from mRNA of five control, five DCM and two ICM heart tissues as indicated. Experiments were normalized

by adjusting the overall signal intensity of each hybridization to 100%, relative expression levels are given.

Fig. 9 a shows the cDNA sequence of clone 66214 (corresponds to SEQ ID NO: 27).

5

Fig. 9 b shows the sequence of the EST clone 66214cds (corresponds to SEQ ID NO: 18). The poly(A) signal is underlined, start and stop codons are marked by bold letters, the sequence of 66214 is marked in italic letters.

10 Fig. 9 c shows the putative amino acid sequence 66214pep (corresponds to SEQ ID NO: 8).

Fig. 9 d shows a schematic alignment of the cDNA fragment 66214 identified in SSH with the Genbank entree and open reading frame of 88 amino acids (aa). Not to scale. Homology scores were determined using blast2 algorithm of NCBI: 66214-AF129505: Expect = e-157, Identities = 290/290 (100%), Positives = 290/290 (100%).

15

Fig. 9 e: Two filters were hybridized sequentially with [ $\alpha$ -33P]UTP labeled T3 transcripts from cDNA libraries prepared from mRNA of five control and five DCM heart tissues as indicated. Experiments were normalized by adjusting the overall signal intensity of each hybridization to 100%, relative expression levels are given. NF1 was not taken into account for calculation of mean values and standard deviations.

20

25

Fig. 10 a shows the cDNA sequence of clone 66268 (corresponds to SEQ ID NO: 28), 52474 (corresponds to SEQ ID NO: 29) and S1MC01-1 (corresponds to SEQ ID NO: 30).

Fig. 10 b shows the sequence of the EST clone X83703 (corresponds to SEQ ID NO: 19). Start and stop codons are marked by bold letters, the sequences of 66268 and SIMC01-1 are marked in italic letters. Multiple AU-rich mRNA decay elements are present in the 3'-noncoding region (underlined).

Fig. 10 c shows the putative amino acid sequence CAA58676 (corresponds to SEQ ID NO: 9). Amino acids 94-97 of 66268 encode a nuclear localization signal pattern 4 ([KR][KR][KR][KR]). The protein is described to be located in the nucleus. Moreover, a PEST-rich region (aa 108-126), a tyrosine phosphorylation site (aa 33) and a domain containing four tandem ankyrin-like repeats (aa 152-183) have also been found.

Fig. 10 d shows a schematic alignment of the cDNA fragments identified in SSH and FDD, respectively with their homologous Genbank entree and the open reading frame of 319 amino acids (aa). Not to scale. Homology scores were determined using blast2 algorithm of NCBI:

66268-X83703: Expect = 9e-77, Identities = 152/152 (100%), Positives = 152/152 (100%)

52474-X83703: Expect = 6e-23, Identities = 59/59 (100%), Positives = 59/59 (100%)

SIMC01-1-X83703: Expect = e-115, Identities = 227/234 (97%), Positives = 227/234 (97%).

Fig. 10 e shows RNA samples prepared from three control, four DCM, three ICM and one HCM heart tissue have been compared by fluorescence differential display using the primer combination [T7]T12MC and [M13r]ARP1 (with the arbitrary sequence CGACTCCAAG). The relative expression was calculated using ImageQuant Software and the lowest value set to 1 as reference for all values. Mean values and standard deviations were calculated from all NF and DCM samples, as well as from ICM75 and ICM96.



Fig. 10 f depicts the recombinant over expression of a 66268-YFP fusions protein in pCMs. The pCMs were transfected with an expression plasmid for a 66268-YFP fusions protein and stimulated with Phenylephrine (100  $\mu$ M). The YFP signal was detected with a fluorescence microscope (Axiovert 100S, Zeiss (Jena); YFP filter set, AF-Analysetechnik (Tübingen)) in combination with a digital camera (LAS-1000, Fuji; AIDA-software, Raytest).

## Examples

The following examples illustrate the invention. These examples should not be construed as limiting: the examples are included for purposes of illustration and the present invention is limited only by the claims.

### EXAMPLE 1

#### 1. Isolation of total RNA from heart tissue

Total RNA was isolated from tissue of explanted hearts of left ventricle of human non-failing and DCM patients, which are listed in TABLE 1, respectively, according to the protocol of Chomczynski and Sacchi with some minor modifications. 0.5 g tissue were disrupted using a mortar and pestle and grinded under liquid nitrogen. The suspension of tissue powder and liquid nitrogen was decanted into a cooled 50 ml polypropylene tube and nitrogen allowed to evaporate completely without thawing the sample. After addition of 10 ml solution D (4 M guanidinium thiocyanate, 25 mM sodium citrate pH 7.0, 0.5 % sodium-N-lauroyl-sarcosinat, 0.1 M 2-mercaptoethanol) the sample was homogenized immediately using a rotor-stator homogenizer (Ultra-Turrax T8, IKA Labortechnik) for 60 s at maximum speed. The sample was mixed with 1 ml 2 M NaOAc pH 4.0, 10 ml phenol (water saturated, pH 4.5-5) and 2 ml chloroform/isoamylalcohol (49/1). After incubation on ice for 15 min and centrifugation at 10000g for 30 min at 4 °C the aqueous phase was transferred to a fresh 50 ml polypropylene tube. RNA was precipitated with 1 vol isopropanol at -20 °C for at least one hour. After centrifugation at 10000g for 30 min at 4

°C the RNA pellet was redissolved in 5 ml Solution D and precipitated again with 1 vol isopropanol as described. The pellet was washed with cold 75% EtOH and dried at RT for 15 min. To completely dissolve RNA 500 µl DEPC-treated water were added and the sample was incubated at 60 °C for 10 min, final storage was at -80 °C. An aliquot was used  
5 for quantification by A<sub>260</sub> measurement and separation on a formaldehyde agarose gel (Sambrook *et al.*) to check integrity and size distribution.

TABLE 1: Human heart samples

ID heart	ID library	diagnosis	sex	age	medication	explantation date
Normal controls						
GH1	NF9	cerebral hemorrhage	f	53	unknown	18.05.1995
GH2	NF19	unknown	m		unknown	
h92	NF1	(suspicion on hepatitis B)	f	50	unknown	20.07.1994
KN3	NF45	intracranial pressure at astrocytoma IV	f	41	Minirin, Dopamin, Rocephin, Dexamethason	30.08.1996
KN4	NF46	traumatic brain disease	m	33	Arterenol, KCl	08.06.1997
KN6	-	unknown			unknown	06.07.1997
KN7	-	unknown			unknown	02.01.1998
DCM samples						
h43	DCM6	DCM	f	54	Digitalis, diuretics, nitrates, ACEI	24.04.1990
h44	DCM13	DCM, Z.n. myocarditis	m	22	unknown	08.05.1990
h94	DCM15	DCM	m	16	Digitalis, ACEI, nitrate, catecholamines, diuretics	03.11.1994
h97	DCM14	DCM	m	62	Digitalis, diuretics, ACEI, Amiodaron, Marcumar	04.01.1995
h99	DCM49	DCM	m	64	Digitalis, diuretics, ACEI, Amiodaron, Marcumar, nitrate	17.05.1995
h100	-	DCM			unknown	20.09.1996
DH2M1	-	DCM	m	53	unknown	
ICM samples						
h75	-	ICM			unknown	05.10.1992
h79	-	ICM			unknown	20.04.1993
h80	ICM47	ICM			unknown	10.06.1993
h81	ICM48	ICM			unknown	17.06.1993
h96	-	ICM	m	39	Digitalis, ACEI, Amiodaron, Marcumar	13.12.1994
HCM samples						
h48	-	non-obstructive HCM	m	37	unknown	08.01.1991

## 2. Isolation of poly(A) RNA from total RNA

Poly(A) RNA was isolated from 300 µg total RNA (see 1.) using the PolyA Quick mRNA Isolation Kit (Stratagene) according to the manufacturers protocol. Purified mRNA was dissolved in 30 µl RNase-free water (Stratagene), quantified and analyzed on a formaldehyde agarose gel as described (see 1.).

## 3. Suppression subtractive hybridization (SSH)

### 3.1 Construction of a subtracted library

2 µg of tester mRNA and 2 µg of driver mRNA were used to construct a subtracted and normalized cDNA library using the PCR-Select cDNA Subtraction Kit and Advantage cDNA-Polymerase Mix (Clontech) according to the manufacturers protocol. In general, two libraries were constructed for each tester and driver combination, since only transcripts can be identified that are over-represented in the tester mRNA.

Both, the subtracted and non-subtracted cDNA population were analyzed on an agarose gel as described (Clontech) and transferred onto Zeta-Probe GT nylon membrane (BioRad) by capillary forces (Sambrook *et al.*). The membrane was UV crosslinked in a Stratalinker 2400 (Stratagene).

To analyze the subtraction efficiency the membrane was hybridized with a Digoxigenin-labeled probe synthesized from a housekeeping gene using the Dig-DNA Labeling and Detection Kit (Roche). For probe synthesis a 451 bp fragment of human GAPDH was amplified from 0.5-1 µg cDNA of a NF heart library (see 5.1.) in a 100 µl PCR reaction with the primer pair provided by the PCR-Select cDNA Subtraction Kit (Clontech). 100 ng of gel purified (QIAquick Gel Extraction Kit, Qiagen) GAPDH cDNA fragment then were used for Dig-labeling. The hybridized membrane was exposed to a X-ray film (X OMAT AR, Kodak) for 15 min. Only subtractions, where the GAPDH signal intensity of the subtracted cDNA population was at least four fold lowered compared to the corresponding non-subtracted cDNA-population, were selected for further analysis. 17 µl of the subtracted sample were purified using a PCR Purification Kit (Qiagen) and eluted in 20 µl ddH<sub>2</sub>O (Gibco BRL).

For addition of 3'-A overhangs 15.7 µl of the purified subtracted cDNA sample was incubated in the presence of PCR buffer, 1.5 U Taq DNA polymerase (APB) and 0.2 mM

dATP for 11 min at 72 °C. 3 µl of the sample was ligated into the pGEM-T easy vector (Promega) and competent *E. coli* cells were transformed as described by the manufacturer.

### 3.2 Amplification of subtracted cDNA clones

Subtracted cDNA clones were grown over night at 37 °C in 96 well microplates filled with  
5 100 µl LB medium (Sambrook *et al.*) and supplemented with 10 µg/ml Amp. 1 µl of the bacterial culture then was transferred into 99 µl PCR premix (1x PCR buffer, 2.5 U Taq DNA polymerase (APB), 0.2 mM dNTP) and directly amplified using the nested primer pair 1 and 2R provided by the PCR-Select cDNA Subtraction Kit (Clontech). Best results were obtained with 27 cycles and an annealing and polymerization temperature of 68 °C.  
10 The size distribution of PCR-products was analyzed on an 1% agarose gel (Sambrook *et al.*). Bacterial cultures were mixed with glycerol to a final concentration of 20% and stored at -80 °C.

## 4. Fluorescence differential display (FDD)

### 15 4.1 DNaseI digestion

Total RNA (see 1.) was digested using the MessageClean-Kit (GeneHunter) according to the manufacturers protocol.

### 4.2 Reverse transcription

Four degenerated primer pools [T7]-T<sub>12</sub>MA, [T7]-T<sub>12</sub>MC, [T7]-T<sub>12</sub>MG and [T7]-T<sub>12</sub>MT  
20 anchoring to poly(A) tails of mRNAs were used, where M is the degenerated position (a mixture of A, C, G). A 17 nt T7 RNA polymerase promoter-derived site (ACGACTCACTATAGGGC) is incorporated which allows the generation of an antisense transcript. For each RNA sample four separate reactions were performed.

200 ng of DNA-free RNA (see 4.1.) was denatured for 5 min at 70 °C in the presence of 0.2  
25 µM anchor primer [T7]-T<sub>12</sub>MX and 20 U rRNasin (Promega). After addition of RT buffer (Gibco), 10 mM DTT, 25 µM dNTP and 200 U SuperscriptII RTaseII (Gibco) on ice, the reaction with a final volume of 20 µl was performed for 5 min at 42 °C and 1 h at 50 °C. The reaction was stopped by heating 15 min at 70 °C.

### 4.3 PCR

Resulting cDNAs (see 4.2.) were reamplified in the presence of the same anchor primer labeled with Cy5 and a second primer with 10 nt of arbitrary chosen sequence. A 16 nt segment of the M13 universal reverse (-48) 24mer priming sequence (ACAATTTTCACACAGCA) is incorporated in the arbitrary primer [M13r]-ARPX<sub>10</sub> for direct sequencing.

1 µl of reverse transcription sample (see 4.2.) was mixed on ice with 1x PCR buffer (Qiagen), 3.75 mM MgCl<sub>2</sub>, 0.35 µM Cy5-[T7]-T<sub>12</sub>MX, 0.35 µM [M13r]-ARPX<sub>10</sub>, 50 µM dNTP and 0.5 U Taq polymerase (Qiagen) in a final volume of 20 µl. PCR was run in a Peltier Thermal Cycler PTC 200 (MJ Research) under the following conditions: 2 min 95 °C, [15 s 92 °C, 30 s 50 °C, 2 min 72 °C]<sub>4</sub>, [15 s 92 °C, 30 s 60 °C, 2 min 72 °C]<sub>25</sub>, 7 min 72 °C, 4 °C.

### 4.4 Electrophoresis on a 6% denaturing polyacrylamide gel

The PCR sample (20 µl, see 4.3.) was mixed with 6 µl gel loading dye (95% formamide, 20 mM EDTA, 0.005% BPB), denatured for 2 min at 80 °C and separated on a standard sequencing gel (6% polyacrylamide/8.3 M urea) at 55 W for 3 h. The gel was dried on Whatman 3MM paper and fluorescence signals read at 635 nm on a Storm fluorimager (Molecular Dynamics). Data analysis was performed using ImageQuant Software (Molecular Dynamics) as described below (see 6.3.).

### 4.5 Recovery of PCR fragments from the sequencing gel

Individual bands of interest (see 4.4.) were cut out of the gel with a scalpel. The gel slice attached to Whatman paper was soaked for 1 h at 37 °C (300 rpm) in 100 µl buffer EB (Qiagen) and incubated at 4 °C over night. Supernatant was purified using the QIAquick PCR purification Kit (Qiagen) as described by the manufacturer. DNA was eluted into 30 µl EB buffer (Qiagen).

### 4.6 Reamplification of differential display PCR fragments

All PCR fragments recovered from the differential display gel could be reamplified with a set of universal primers, M13r(-48) primer [AGCGGATAACAATTTTCACACAGGA] and T7 primer [GTAATACGACTCACTATAGGGC]. A 40 µl PCR was set up on ice with 3 µl template (see 4.5.), 1x PCR buffer, 1.5 mM MgCl<sub>2</sub>, 20 µM dNTP, 0.2 µM T7 primer, 0.2

$\mu$ M M13r(-48) primer and 2 U Taq polymerase (Qiagen) and run as described above (see 4.3.).

#### 4.7 Electrophoresis on a preparative 1.2% agarose gel

30  $\mu$ l of reamplified PCR sample were mixed with 6  $\mu$ l loading dye and separated on an  
5 1.2% agarose/1x TBE gel together with a size standard and a PCR marker (Promega). Bands were cut out with a scalpel and DNA extracted from agarose gel slice using QIAquick gel extraction Kit as described (Qiagen). 1  $\mu$ l of recovered DNA was used for sequencing.

### 10 5. Preparation of cDNA libraries and probe synthesis

Since the availability of heart material is very limiting, labeled *in vitro* transcripts of a cDNA library prepared from heart mRNA were used for dot blot hybridization instead of reverse transcribed mRNA itself.

#### 5.1 Preparation of a cDNA library

15 5  $\mu$ g of high quality mRNA (see 1., see 2.) were used to prepare a cDNA library using the cDNA Synthesis Kit and ZAP-cDNA Gigapack III Gold Cloning Kit (Stratagene) as described in the manual with the following modifications:

(a) Packaging and titering: 2.5  $\mu$ l of the ligation reaction were packaged. If the library did not represent at least one million clones, the remaining 2.5  $\mu$ l were also packaged.  
20 After centrifugation of XL1-Blue MRF' culture (50 ml), the cells were gently resuspended in 10 mM MgSO<sub>4</sub> at 4 °C and immediately used for transduction or stored for max 40 h at 4 °C.

(b) Determination of the insert size: 25 plaques were transferred from agar plates used for titering directly into 40  $\mu$ l PCR premix (1x PCR-buffer, 0.25  $\mu$ M T3 primer, 0.25  
25  $\mu$ M T7 primer, 200  $\mu$ M dNTP, 0.085 U Taq DNA-polymerase) and inserts amplified using 35 cycles and an annealing temperature of 48 °C. The insert size was checked on an agarose gel and was in the range of 1-2 kb.

(c) Storage of the library: Libraries were transferred into 50 ml-polypropylene tubes, supplemented with 150  $\mu$ l 0.3 % chloroform and stored at 4 °C. A part of each library  
30 was stored in 7 % DMSO at -80 °C.

Mass *in vivo*-excision was done according to the protocol of the ZAP-cDNA Gigapack III Gold Cloning Kit with the following modifications:

Transfected XL1 Blue MRF' were grown in 25 ml LB. 5 ml of the supernatant containing single stranded phages was used to infect 20 ml of SOLR cells. Remaining 20 ml of single stranded phages were stored at 4 °C for up to two months. To determine the titer of excised phagemids 10 µl, 1 µl and 0.1 µl of infected SOLR cells were plated on LB/Amp dishes. If the titer was lower than one million, 5 ml or more of the remaining supernatant was used again to infect fresh SOLR cells. Infected SOLR cells (25 ml) were grown in 200 ml LB/Amp over night for plasmid isolation (Plasmid Midi Kit, Qiagen).

#### 5.2 Linearization of the template cDNA library for *in vitro* transcription

200 µg plasmid DNA were digested with XhoI over night at 37 °C in a volume of 250 µl to linearize the plasmid at the 3' end of the insert. The sample was controlled for complete digestion on an agarose gel, treated with 10 µg/µl Proteinase K (Roche) at 37 °C for 30 min, extracted once with TE saturated phenol (pH 7.5-8) and once with chloroform/isoamylalcohol (24/1) and precipitated in the presence of 0.1 volume 3 M NaOAc (pH 5.2) and 3 volume EtOH. The pellet was washed with 500 µl 75% ethanol, dried at RT for 10 min, dissolved in 150 µl DEPC-treated water and quantified.

1 µg of linearized plasmid was used for an *in vitro* transcription as described (see 5.3.), omitting the radioactive labeled nucleotide and adding UTP to a final concentration of 10 mM. Following DNaseI digestion, the RNA was extracted with phenol/chloroform/isoamylalcohol (24/23/1), precipitated with EtOH and dissolved in 15 µl DEPC-treated water. The yield was in the range of 15-22 µg RNA. 1.5 µl RNA were separated on a formaldehyde agarose gel. A smear of transcripts was visible between 0.5 kb and 10 kb with a peak at about 1 kb.

#### 5.3 *In vitro* transcription

According to the RNA Transcription Kit (Stratagene) 1 µg of linearized template (see 5.2.) was incubated in the presence of 1x transcription buffer, 10 mM ATP, 10 mM CTP, 10 mM GTP, 1 mM UTP, 70 µCi [ $\alpha$ -<sup>33</sup>P]UTP (APB), 0.75 M DTT, 20 U rRNasin (Promega) and 25 U T3 RNA polymerase for 30 min at 37 °C. After addition of 5 U RNase-free DNaseI (Roche) the sample was incubated for 15 min at 37 °C. 25 µl STE-buffer (APB) was added



to the probe and the reaction purified using G50 Micro Columns (APB) according to the manufacturers protocol.

#### 5.4 Prehybridization of *in vitro* transcripts

To suppress probe hybridization to human repetitive DNA, labeled RNA was prehybridized  
5 to cot1-DNA. 213 µl DEPC-treated water, 100 µl 20x SSC, 2 µl 20% SDS and 40 µl cot1-DNA (1 µg/µl, Gibco BRL) were added to 45 µl labeled RNA (see 5.3.). denatured at 95 °C for 2 min and incubated for 2 h at 65 °C.

### 6 Quantitative Dot Blot Analysis

#### 10 6.1 Transfer of PCR fragments onto nylon membrane

For spotting, approximately 300 ng PCR product (see 3.2.) or gene-specific control cDNA fragments were mixed with 140 µl 0.4 M NaOH/10 mM EDTA pH 8.0 in 96 well microplates and denatured 10 min at 95 °C. 50 µl of each PCR-fragment (at least 100 ng cDNA) were transferred on a nylon membrane (11.4x7.5 cm. BioRad) using a 384 hole  
15 vacuum apparatus (Keutz, custom-made). 50 µl 0.4 M NaOH were added to each position and transferred. The membrane was washed in 2x SSC, dried for at least 1 h at RT and fixed by UV crosslinking (Stratalinker 2400, Stratagene). For each experiment two identical membranes were prepared in parallel.

#### 6.2 Dot blot hybridization and washing

20 The cDNA filter was soaked in 2x SSC and transferred into a hybridization flask. The membrane was hybridized with 10 ml hybridization solution (6x SSC, 5x Denhardts, 0.2 % SDS, 0.2 % sodium pyrophosphate) supplemented with 50 µg/ml denatured salmon sperm DNA (Typ III, Sigma) at 65 °C for 2 h in an Unitherm 6/12 hybridization oven (UniEquip). The prehybridization mix was poured off. 200-400 µl of cot1-hybridized probe (see 5.4.)  
25 were added to 8 ml of hybridization solution (including salmon sperm DNA) preheated to 65 °C. Dot blots were hybridized over night at 65 °C. For washing of cDNA filters all solutions were heated to 65 °C. The membrane was washed twice with 50 ml wash solution 1 (2x SSC, 0.1 % SDS) for 30 min, then twice with 50 ml wash solution 2 (0.1x SSC, 0.1 % SDS) for 30 min and wrapped in a keep-fresh foil. The filter was exposed to a phosphor

screen for two days and scanned at 450 nm using the Storm Phosphoimager (Molecular Dynamics).

### 6.3 Data analysis

Signal intensities were calculated using ImageQuant Software (Molecular Dynamics) by subtracting the local background. For comparison of different filters signal intensities were normalized by adjusting the overall intensity of each filter to 100%. In general, two cDNA filters were hybridized successively with 10 probes prepared from different human heart samples.

Dots which represented at least two fold changes in signal intensity comparing the group of DCM heart samples (y) with that of normal controls (x) were selected for further analysis. The probability of type 1 error was calculated to be less than 5% using the Wilcoxon test. This non-parametric statistic algorithm does not assume any distribution of x and y values. If the sample size of one group was smaller than 4 the Wilcoxon test could not be applied. Instead significance of gene regulation was confirmed by a t-test. The t-test assumes that standard deviations of both groups x and y are similar and values distributed according to normal distribution.

Independent of the disease individual differences between human samples are expected. They are the result of the different genetic background of individuals, sex, age, environmental and life conditions (e.g. smoking, drinking, nourishment), the status of disease and medical treatment. Especially DCM patients were treated by a number of drugs prior to heart transplantation. We laid down that the regulation has to be consistent in at least two DCM patients and more or less homogenous in all but one non-failing patient.

Selected clones were grown in 5 ml LB/Amp from glycerol stocks (see 3.2.). Plasmids were isolated using the Plasmid Mini Kit (Qiagen) and sequenced.

### 6.4 Stripping of dot blot membranes

cDNA filters were transferred into boiling stripping solution (0.1x SSC, 0.5 % SDS) and incubated for 1 h at RT. This procedure was repeated until no more radioactivity could be detected by a Geiger-Müller counter. The filter again was wrapped in keep-fresh foil and stored at RT.

## 7. Full-length cloning:

Full-length cloning was performed using RT-PCR with oligonucleotides priming to the 5'- and 3'- ends of the sequence encoding the open reading frame. PCR-fragments were then purified by agarose gel-electrophoresis followed by gel elution using the gel purification kit from Qiagen. PCR-fragments were finally cloned into p201-DONOR (Life Technologies) or pTOPO2.1 (Invitrogen).

The cloned cDNAs were verified by sequencing. In addition, *in vitro* translations were performed using the TNT Quick Coupled Transcription/Translation Systems (Promega) in order to verify the correct molecular weight of the proteins encoded by a given cDNA. The full-length clones were named according to their ID number provided with the suffix "-cds" (xxxxx-cds). The proteins were named according to their ID number provided with the suffix "-pep" (xxxxx-pep).

## 8. Yeast two-hybrid system

### 8.1 Two-hybrid screen protocol (Golemis et al., 1994).

The yeast two-hybrid vectors are described in section below. Yeast strains used were EGY48LacZ-GFP (*ura3::6\*LexOp-lacZ*, *lys2::6\*LexOpCYC1GFP*, *his3*, *trp1*, *6\*LexAOp-LEU2*, *mat $\alpha$* ) and EGY199UL (*ura3::6\*LexOp-lacZ*, *his3*, *trp1*, *6\*LexAOp-LEU2*, *mat a*). Yeast was grown in YPD or selective minimal medium (Sherman 1986). Transformations were done using the high-efficiency method of Gietz et al., 1992. The bait plasmids were first introduced in the yeast strain EGY48LacZ-GFP resulting in the strain EGY48LacZ-GFP-bait. Self activation of the bait was checked by plating the yeast on minimal glucose medium with or without X-Gal (5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside). In parallel protein expression was verified by western blot analysis using a polyclonal rabbit anti-LexA antiserum. A human heart cDNA library (pJG#19) cloned (EcoRI/XhoI) in the vector pJG4-5 was then introduced in the EGY48LacZ-GFP-bait strain. After transformation  $4 \times 10^4$  colonies per plate) yeast were plated on selective medium (-histidine, -tryptophane, +methionine, glucose). Colonies were harvested and an aliquot was plated on selective medium (-histidine, -tryptophane, -uracil, raffinose, galactose, X-gal). The interactions were assayed by colony growth on selective medium as

well as by  $\beta$ -galactosidase activity on the plate. Positive clones were plated over night on medium (-histidine, -tryptophane, -uracil, glucose, X-gal) in order to deactivate the expression of the prey. The verification of the interaction was performed by plating the colonies on medium A:(-histidine, -tryptophane, -uracil, glucose, X-gal) and medium B: (-histidine, -tryptophane, -uracil, raffinose, galactose, X-gal). Only blue colonies growing on medium B but not on medium A were further analysed by yeast-colony-PCR. Plasmids were rescued and introduced in *E.coli* (Robzyk and Kassir, 1992). DNA was isolated from the bacteria and sequenced. Interactions were finally verified by reintroducing the plasmid (prey) in the yeast strain EGY199UL. Mating of the EGY199UL (prey) with the corresponding EGY48LacZ-GFP (bait) was performed in order to get a diploid strain carrying bait and prey (Guthrie and Fink, 1991; Pringle et al., 1997; Golemis and Khazak, 1997). Protein interaction resulted in growth and blue colour of the diploid colonies on medium B but not on medium A. Interactions were further analysed by quantifying the relative activity of the GFP reporter in a FACS assay.

## 8.2 Two hybrid vectors description

### 8.2.1 Bait vectors

- 1) pSH2-1 (Hanes SD. and Brent R. 1989)
- 2) pEG202(U8996)
- 3) 413MetLexN0

The vector 413MetLexN0 was constructed by cloning a PCR generated full length LexA repressor cDNA (with XbaI/BamHI overhangs) into the vector 413Met25 (Mumberg et al., 1994) cut XbaI/BamHI.

- 4) 413MetLexN0.att

The destination vector 413MetLexN0.att was constructed by introducing the rfC cassette of the Gateway<sup>TM</sup> system (Invitrogen) into the vector 413MetLexN0. For this purpose a linear PCR fragment comprising the rfC-cassette and flanking homologies of 40 bp to the LexA gene or 40 bp(5-prime) of the CYC1 terminator(3-prime) of the vector 413MetLexN0 was used for homologous recombination to the EcoRI linearized vector 413MetLexN0 in yeast. One correct recombinant vectors was re isolated from yeast and can be used for cloning of cDNAs by in vitro recombination performing a LR-reaction of the Gateway<sup>TM</sup> system.

## 5) 413MetLexC0

The vector 413MetLexC0 was constructed by cloning a PCR generated full length LexA repressor cDNA (with HindIII-ClaI-XhoI/Sall overhangs) into the vector 413Met25 (Mumberg D et al., 1994) cut HindIII/XhoI.

## 5 6) 413MetLexC0.att

The destination vector 413MetLexC0.att was constructed analogous to the procedure described for the vector 413MetLexCN.att.

**8.2.2 Prey vectors**

## 1) pJG4-5(U89961)

## 10 2) 424GBN0

The vector 424GBN0 was constructed by cloning a PCR generated full length B42 transactivation domain cDNA (with XbaI/BamHI overhangs) derived from the vector pJG4-5 into the vector 424GAL1(Mumberg D et al., 1994) cut SpeI/BamHI.

## 3) 424GBN0.att

15 The destination vector 424GBN0.att was constructed by introducing the rfC cassette of the Gateway™ system (Invitrogen) into the vector 424GBN0. For this purpose a linear PCR fragment comprising the rfC-cassette and flanking homologies of 40 bp to the LexA gene or 40 bp(5-prime) of the CYC1 terminator(3-prime) of the vector 424GBN0 was used for homologous recombination to the EcoRI linearized vector 424GBN0 in yeast. One correct  
20 recombinant vector was re-isolated from yeast and can be used for cloning of cDNAs by in vitro recombination performing a LR-reaction of the Gateway™ system.

## 4) 424GBC0

The vector 424GBC0 was constructed by cloning a PCR generated full length B42 transactivation domain cDNA (with HindIII-ClaI-XhoI/Sall overhangs) into the vector  
25 424GAL1 (Mumberg D et al., 1994) cut HindIII/XhoI.

## 5) 424GBC0.att

The destination vector 424GBC0.att was constructed analogous to the procedure described for the vector 424GBCN.att.

**8.3 Two-hybrid interaction matrix (40K matrix)**

30 A collection of yeast two-hybrid 200 plasmids (baits and preys) made at Medigene was introduced in EGY48LacZ-GFP and EGY199UL respectively. Each EGY48LacZ-GFP-bait

were challenged against each EGY199UL-prey for interaction via mating (Golemis and Khazak, 1997). The resulting interactions tested were  $40.10^3$ . This procedure correspond to the MediGene 40K matrix. Positive interaction were scored by growth on selective medium and  $\beta$ -galactosidase activity. Moreover, the strength of the interactions were quantified in a  
5 FACS assay. All interactions were stored in the programme CACI (Computer analysis of Complex Interactions). Matrix interaction analysis was performed using the programme CACI.

## 9. Recombinant gene expression in cardiomyocytes

### 10 9.1 Isolation of primary cardiomyocytes from neonatal rats

Neonatal rats (P2-P7) were sacrificed by cervical dislocation. The ventricles of the beating hearts were removed and cardiomyocytes were isolated with the "Neonatal Cardiomyocyte Isolation System" (Worthington Biochemicals Corporation, Lakewood, New Jersey) according to the protocol. Briefly, the ventricles were washed twice with ice cold Hank's  
15 Balanced Salt Solution without Potassium and Magnesium (CMF-HBBS) and minced with a scalpel to an average volume of one cubic millimeter. The heart tissue was further digested over night with trypsin at  $10^{\circ}\text{C}$ . Next morning trypsin inhibitor and collagenase were added. After an incubation at  $37^{\circ}\text{C}$  and mild agitation for 45 minutes the cells were dispersed by pipetting. The solution was further purified by  $70\text{ }\mu\text{m}$  mesh (Cell Strainer) and  
20 centrifuged twice for 5 minutes at  $60 \times g$ . The cell pellet was resuspended in plating medium and counted. Cells were seeded with a density of  $2 \times 10^4/\text{cm}^2$  on gelatine (Sigma, Deisenhofen) coated dishes. The next morning cells were washed twice with DMEM and maintenance medium was added.

25 Plating medium: DMEM/M-199 (4/1); 10% Horse serum, 5% Fetal calf serum;  
1 mM sodiumpyruvate; antibiotics and antimycotics  
Maintenance medium: DMEM/M-199 (4/1); 1 mM sodiumpyruvate

### 9.2 Construction of expression plasmids for cardiomyocytes

30 The pCI-vector (Promega) was cut with BsrGI. The linearized vector was incubated with the Klenow-fragment and dNTPs to generate blunt ends. The resulting vector was cut with

NheI and NotI after religation and gel purified. A PCR fragment comprising the entire open reading frame without the start codon of the yellow variant of the green fluorescent protein (YFP) was inserted into the NheI and NotI sites. The PCR was performed under standard conditions with the following primers to add several unique restriction site for further cloning:

5'-primer: SpeI-XbaI-EcoRI-XhoI-YFP

5'-GGA CTA GTT CTA GAG AAT TCC TCG AGG TGA GCA AGG  
GCG AGG AG-3'

3'-primer: YFP-STOP-NotI (the NotI site was derived from the vector)

5'-AGT TGG TAA TGG TAG CGA CC-3'

template: pEYFP-vector (Clontech)

The PCR product was gel purified and digested with SpeI and NotI to generate compatible ends. The resulting vector was linearized with XbaI and EcoRI and gel purified in order to insert a consensus Kozak-sequence, which was derived from oligo annealing.

5'-Kozak: 5'-CTA GAA CTA GTT CCA CCA TGG-3'

3'-Kozak 5'-AAT TCC ATG GTG GAA CTA GTT-3'

In the final construction step the plasmid was linearized with EcoRI and XhoI and gel purified. A PCR fragment comprising the entire open reading frame of 66268 flanked by an EcoRI site at the 5'-end and a XhoI site at the 3'-end was inserted.

### 9.3 Stimulation of isolated cardiomyocytes from neonatal rats

Stimulation of primary cardiomyocytes from neonatal rats (pCMs) was started two to six hours after medium was changed to maintenance medium. Directly after stimulation pCMs were infected with recombinant adenoviruses at a MOI of five. Cells were incubated for 48 hours at humidified atmosphere at 37°C and 5% CO<sub>2</sub> followed by an analysis of morphological alterations.

### 9.4 Transient transfection of isolated cardiomyocytes from neonatal rats

For each well of a six well plate 1 µg of plasmid DNA was combined with 20 µl 2 x BBS and 100 µl maintenance medium without antibiotics. Meanwhile 4 µl of LIPOFECTAMINE (Gibco/BRL) were mixed with 650 µl maintenance medium without antibiotics in a polystyrene tube. The DNA-sample was added after an incubation for 15' at room-temperature. The suspension was mixed by inverting the tube twice and incubated for

15' at room-temperature. Meanwhile medium was changed to 1 ml maintenance medium without antibiotics. The transfection-mixture was added onto the cells and gene expression was analysed 48 hrs later.

2 x BBS:                50 mM BES  
5                        280 mM NaCl  
                         1.5 mM Na<sub>2</sub>HPO<sub>4</sub>  
                         adjust to pH 6.95 by administration of NaOH

## EXAMPLE 2

10 EST 40399 (FIG. 1A) was identified by suppression subtractive hybridization comparing transcript levels of heart tissue explanted from normal control h92 with one from DCM patient h97 (see TABLE 1). The fragment was found to be over-represented in the DCM tissue.

15 As of FIG. 1 D the identified cDNA fragment is a part of the EST clone NM\_003970 (FIG. 1 B), which encodes the amino acid sequence NP\_003961 (identical to CAA48832; FIG. 1 C). This amino acid sequence encodes the 165 kDa M-protein, also known as myomesin 2 or MYOM2.

20 Z and M bands of the sarcomere are interconnected by the long titin molecules. The 165 kDa M-protein is one of two known titin-associated proteins, which seem responsible for the formation of a head structure on one end of the 0.9 micron long titin string (Vinkemeier *et al.*). M-protein may function in strengthening the links between thick filaments necessary to withstand the stronger tension during contraction in the heart and in fast fibers (van der Ven *et al.*)

25 Upregulation upon DCM was confirmed for two additional DCM patients compared to five normal control hearts by quantitative dot blot analysis (FIG. 1 E). The relative expression level of 40399 is induced by a factor of 3.1 upon disease. The probability of type 1 error is less than 5% as determined in a t-test.

30 Expression was not induced in two DCM patients, which may reflect individual differences throughout the population.



Significant upregulation of 40399 expression in heart tissue of two DCM patients compared to five normal controls indicates that an increased expression of 40399 is associated with dilated cardiomyopathy. Upregulation of titin-associated muscle M-protein by a factor of 3 may massively interfere with normal myofibril assembly and stabilization and decrease muscular activity. From our data we conclude that abnormalities in expression of this protein are associated with muscular abnormalities that result in cardiomyopathies. Therefore we expect the protein to play a causative role in heart diseases, especially congestive heart failure.

Mutations in other sarcomeric proteins have already been identified as causes of hypertrophic cardiomyopathy, suggesting that cytoskeletal proteins play a central role in cardiac function (Hein *et al.*). These findings support our general observation of a causative correlation between deregulation of sarcomeric proteins and reduced contractile function in end-stage heart failure. Therefore, 40399 can serve as a heart disease marker and a specific molecular target for drug development.

Downregulation of protein expression by specific inhibitors or antisense constructs seems to be a very promising therapeutic tool to treat heart diseases.

### EXAMPLE 3

EST 41441 (FIG. 2 A) was identified by suppression subtractive hybridization comparing transcript levels of heart tissue explanted from normal control h92 with one from DCM patient h97 (see TABLE 1). The fragment was found to be over-represented in the control tissue. The identified cDNA fragment is a part of the EST clone AW755252 (FIG. 2 B), which predicts an amino acid sequence 41441pep given in FIG. 2 C (schematic alignment FIG. 2 D).

Downregulation upon DCM was confirmed for four DCM patients compared to five normal control hearts by quantitative dot blot analysis. The relative expression level of 41441 is reduced by a factor of 4.5 upon disease (FIG. 2 E). The probability of type I error is less than 5% as determined in a Wilcoxon test.

The EST clone AW755252 (Walker *et al.*) was isolated from a human cardiac muscle expression library and found to be similar to cardiomyopathy associated gene 3 (CMYA3, unpublished).

The LIM sequence motif is a part of the cardiomyopathy associated gene 3.

- 5 The LIM sequence motif was first identified in homeodomain proteins Lin-11, Isl-1 and Mec-3. The LIM domain is a double zinc finger motif that mediates the protein-protein interactions of transcription factors, signaling- and cytoskeleton-associated proteins. There is no evidence, that LIM domains bind DNA directly. Instead, an increasing number of studies implicate LIM domains in protein-protein interactions that regulate development,  
10 cellular differentiation and the cytoskeleton (Bach).

#### **Yeast two-hybrid interactions**

- Interactors with the protein coded by 41441pep were screened using 41441pep as a bait. A large screen was performed using 4 large plates for the library transformation which led to  
15 the analysis of  $2 \times 10^7$  clones. The two hybrid procedure described (protocol 22) led to the identification of 4 different interacting partners. The corresponding cDNAs were identified by homology search using the first 500 nucleotides sequence of the pray clone. The partners are: Hepatitis B virus interacting protein (AF029890), U6 snRNA-associated Sm-like protein LSm8 (AF182294), unknown protein HSPC297 (AF161415) and supervillin  
20 (AF051851).

#### **Hepatitis B virus interacting protein or XIP**

- The identity with Hepatitis B virus interacting protein (AF029890) was found to be 100% over the first 400 amino acids. The homology starts at nucleotide 9 of the AF029890  
25 sequence. The XIP cDNA recognizes a single 0.7 kb transcript in all tissues studied and was particularly abundant in skeletal and cardiac muscles tissues (Melegari *et al.*, 1998). The XIP protein was also found to interact with the hepatitis B virus protein HBx (Melegari *et al.*, 1998). Interestingly, over-expression of the XIP protein prevented wild-type HBx activity on such promoters as well as reduced HBV replication to levels comparable to  
30 those observed with an HBx-minus variant strain (Klein *et al.*, 1999 )

#### U6 snRNA-associated Sm-like protein LSm8

The sequence revealed 100% homology to Homo sapiens U6 snRNA-associated Sm-like protein LSm8 over 400 nucleotides. The homology starts at nucleotide 31 of the AF182294 sequence. The yeast homologue of Lsm8 seems to be play a role, together with Lhp1, as a  
5 molecular chaperone of polymerase III. Lsm8 might be implicated in the very early steps of the U6 snRNP assembly (Panome et al., 1998).

#### Supervillin

10 Homology search using the interactor of clone 41441 led to the identification of supervillin (SVIL) (XM\_011894, AF109135) with 99% identity. Supervillin RNA are expressed ubiquitiously. The human supervillin gene is localized to a single chromosomal locus at 10p11.2 a region that is deleted in some prostate tumours as well as in so tumour cell lines (Pope et al., 1998). The cDNA sequence of this interactor showed identity to supervillin  
15 isoform 2, a membrane associated F-actin binding protein. This protein is also known as archvillin or p205. The identity starts at amino acid 1872 and stops at 1997. Alignment with clones of the database showed that the bait encodes the C-terminal part of the protein supervillin. In this sequence the motif GEL (Gelsolin homology domain) could be identified from amino acid 39 to 138. This domain was also found in  
20 Gelsolin/severin/villin. It is thought to exist both as a intra- and extracellular domain and and may be responsible for Calcium-binding as well as actin-binding. This protein is tightly associated with both actin filaments and plasma membrane specifically in focal adhesion plaques. Over-expression of full-length supervillin in these cells disrupts the integrity of focal adhesion plaques and results in increased levels of F-actin and vinculin. Moreover,  
25 supervillin contains nuclear targeting signals in the centre of the protein which seem to be functional. Therefore supervillin may contribute to cytoarchitecture in the nucleus as well as at the plasma membrane (Wulfkühle et al., 1999).

Significant downregulation of 41441 expression in heart tissue of four DCM patients  
30 compared to five normal controls indicates that a lowered expression of 41441 is associated with dilated cardiomyopathy. Lowered expression of 41441 by a factor of 4-5 seems to

induce a cardiomyopathic phenotype. Therefore we expect the protein to play a causative role in heart diseases, especially congestive heart failure.

The predicted functional domain LIM\_1 also indicates a major role of 41441 in regulation of development, cellular differentiation or the cytoskeleton. From our data together with those from Genbank entree AW755252 we conclude that 41441 is predominantly expressed in cardiac muscle, which supports our idea that 41441 can serve as a marker for heart diseases and a specific molecular target for drug development.

Upregulation of protein expression by gene therapeutic intervention, compensatory molecules or specific activators seems to be a very promising therapeutic tool to treat heart diseases.

#### EXAMPLE 4

EST 52706 (FIG. 3 A) was identified by suppression subtractive hybridization comparing transcript levels of heart tissue explanted from normal control KN2 with one from DCM patient DHZM3 (see TABLE 1). The fragment was found to be over-represented in the diseased tissue.

EST 52706 (FIG. 3 A) was found to be repressed upon disease in screens for expression profiles using suppression subtractive hybridization (?). Transcript levels are significantly downregulated by a factor 27,3 in five DCM patients compared to five normal controls (FIG. 3 B). The probability of type 1 error is less than 5% as determined in a Wilcoxon test. Significant homologies to known sequences from Genbank were not found.

Significant downregulation of 52706 expression in heart tissue of six DCM patients compared to the same number of normal controls indicates that a lowered expression of 52706 is associated with dilated cardiomyopathy. The extreme decrease in expression of 52706 by a factor of 27 seems to induce a cardiomyopathic phenotype. Therefore we expect the protein to play a causative role in heart diseases, especially congestive heart failure. As a conclusion 52706 can serve as a marker for heart diseases and a specific molecular target for drug development.

Upregulation of protein expression by gene therapeutic intervention, compensatory molecules or specific activators may be a therapeutic tool to treat heart diseases.

### EXAMPLE 5

5

EST 56461 (FIG. 4 A) was identified by suppression subtractive hybridization comparing transcript levels of heart tissue explanted from normal control KN5 with one from DCM patient h52 (see TABLE 1). The fragment was found to be over-represented in the DCM tissue.

10 The identified cDNA fragment was found to be overlapping with the EST clone AF077035 (FIG. 4 B), which encodes the amino acid sequence AAD27768 (FIG. 4 D). The predicted amino acid sequence for 56461 is shown in sequence 56461pep (FIG. 4 C).

AF077035 was isolated from CD34(+) hematopoietic stem and progenitor cells (HSPC, Zhou *et al.*). The amino acid sequence of AAD27768 is to 91% identical to one translated  
15 from EST AW785791, which was identified to be specifically expressed in pooled tissues from *Sus scrofa* embryos (Fahrenkrug *et al.*).

Upregulation upon DCM was confirmed for two additional DCM patients compared to five normal control hearts by quantitative dot blot analysis (FIG. 4 E). For these samples,  
20 DCM15 and DCM13, the relative expression level of 56461 is induced by a factor of 5.4. The probability of type 1 error is less than 1% as determined in a t-test.

The remaining three DCM patients did not show a significant change in 56461 expression, which may be the result of individual differences throughout the population.

25 Significant upregulation of 56461 expression in heart tissue of three DCM patients compared to six normal controls indicates that an increased expression of 56461 is associated with dilated cardiomyopathy. Increased expression of 56461 by a factor of 5-6 seems to induce a cardiomyopathic phenotype. Therefore we expect the protein to play a causative role in heart diseases, especially congestive heart failure.

Moreover, the homology to RNA binding domains may indicate a regulatory function for 56461. This finding supports our idea that 56461 can serve as a marker for heart diseases, especially congestive heart failure and a specific molecular target for drug development.

Downregulation of protein expression by specific inhibitors or antisense constructs seems  
5 to be a very promising therapeutic tool to treat heart diseases.

### EXAMPLE 6

EST 61105 (FIG. 5 A) was identified by suppression subtractive hybridization comparing  
10 transcript levels of heart tissue explanted from normal control KN4 with one from DCM patient h94 (see TABLE 1). The fragment was over-represented in the control tissue. The identified cDNA fragment was found to be a part of the EST clone M14780 (FIG. 5 B), which encodes the amino acid sequence AAA52025 (FIG. 5 C; schematic alignment FIG. 5 D). This amino acid sequence encodes the muscle isoform of creatine kinase (creatine  
15 kinase M, Perryman *et al.*), which is one of the important structural and energy metabolism components in skeletal muscle. It catalyzes the reversible transfer of phosphoryl group from creatine phosphate to ADP to form ATP to sustain contractile activity.

Downregulation upon DCM was confirmed for five DCM patients compared to the same  
20 number of normal control hearts by quantitative dot blot analysis (FIG. 5 E). The relative expression level of 61105 is significantly reduced by a factor of 4 upon disease. The probability of type I error is less than 5% as determined in a Wilcoxon test.

#### Yeast two-hybrid interaction

25 The interactors were identified using the 40K matrix of MediGene and analysed by MediGene CACI programme. The following three proteins interact with AAA52025: CapZa (P52907), c-Raf (P04049), FBP (AF049528).

CapZa

CapZ alpha has been localized on Chromosome 1 at position 1p36.13-q23.3. CapZa is an Actin capping protein which bind as heterodimer F-actin at the fast growing end in a Ca<sup>2+</sup> independent manner.

5

FBP11 (Formin binding protein):

Synonyms for FBP are: HYPA, huntingtin-interacting protein (AF049528, AF049524, AF049523) and Fas-ligand associated factor (U70667). FBP11 contains WW motifs that recognize PPXY or PPLP motifs to mediate the interaction (Bedford et al., 1997). Creatine-kinase-M contains a PPXY motif at position 143.

10

c-Raf (isoforme of Raf-1)

c-Raf was localised on chromosome 3 a locus 3p25. This protein belongs to the Ser/Thr family of protein kinase, it contains a zinc-dependent phorbol-ester and DAG binding domain. Moreover, a relationship between c-Raf and Creatine kinase has been shown by other groups in myoblasts (Coolican et al., 1997; Samuel, 1999) and in rhabdomyosarcoma (Ramp et al., 1992).

15

Significant downregulation of 61105 expression in heart tissue of five DCM patients compared to the same number of normal controls indicates that a lowered expression of 61105 is associated with dilated cardiomyopathy. Downregulation of creatine kinase M by a factor of 4 massively decreases the energy reservoir which is necessary to sustain muscle contractility. Therefore we expect the protein to play a causative role in heart diseases, especially congestive heart failure.

20

25

The protein expression was also observed to be deregulated upon canine rapid ventricular pacing, which produces a low output cardiomyopathic state similar to DCM (Heinke *et al.*). Taken together, these results strongly support the notion that energy production is impaired and mitochondrial dysfunction is involved in the development of heart failure. These findings support our general observation of a causative correlation between energy

30

depletion and end-stage heart failure. Therefore, 61105 is a marker and in our opinion also a specific molecular target for drug development.

Upregulation of protein expression by gene therapeutic intervention, compensatory molecules or specific activators seems to be a very promising therapeutic tool to treat heart diseases. In general, increasing the level of available energy sources for muscle contraction by increasing the concentration of free ATP or creatine phosphate would be of great benefit in treating heart failure.

#### EXAMPLE 7

EST 61166 (FIG. 6 A) was identified by suppression subtractive hybridization comparing transcript levels of heart tissue explanted from normal control KN4 with one from DCM patient h94 (see TABLE 1). The fragment was over-represented in the control tissue.

Using LabOnWeb (Compugen) it was possible to assemble 61166contig (FIG. 6 B) that codes for a predicted protein with the amino acid sequence of 61166pep (FIG. 6 C). The assembly of EST is shown in FIG. 6 D with examples of known ESTs (AI 745235, AI 050107, AI 927050)

61166 displays a significant homology to human 65 kDa yes-associated protein YAP65 (NM\_006106, Expect = 2e-84, Identity 57%, Wambutt *et al.*). YAP65 associates *in vitro* with the Src homology domain 3 (SH3) of the Yes proto-oncogene product (yes kinase) and other signaling molecules (Sudol *et al.*). The motif PVKQPPPLAP of human YAP65, which binds to SH3 domains is not conserved in 61166 (amino acids 201-210 marked in italic letters above).

Downregulation upon DCM was confirmed for five DCM patients compared to the same number of normal control hearts by quantitative dot blot analysis (FIG. 6 E). The relative expression level of 61166 is significantly reduced by a factor of 3.9 upon disease. The probability of type 1 error is less than 5% as determined in a Wilcoxon test.

Significant downregulation of 61166 expression in heart tissue of five DCM patients compared to five normal controls indicates that a lowered expression of 61166 is associated with dilated cardiomyopathy. Lowered expression of 61166 by a factor of 4 seems to



induce a cardiomyopathic phenotype. Therefore we expect the protein to play a causative role in heart diseases, especially congestive heart failure.

The high homology to a yes kinase associating protein suggests a central role for 61166 in signal transduction or development. This finding supports our idea that 61166 can be used  
5 as a specific molecular target for drug development and/or diagnostics.

Upregulation of protein expression by gene therapeutic intervention, compensatory molecules or specific activators may be a therapeutic tool to treat heart diseases.

### EXAMPLE 8

10 Screen for expression profiles using a dot blot ybridization in a higher number of patients clearly showed that 61244 is induced upon disease (FIG. 7 E). Transcript levels are significantly upregulated by a factor 3.6 in five DCM patients compared to five normal controls. The probability of type 1 error is less than 5% as determined in a Wilcoxon test.

15 EST 61244 (FIG. 7 A) was identified by suppression subtractive hybridization comparing transcript levels of heart tissue explanted from normal control KN4 with one from DCM patient h94 (see TABLE 1). The fragment was found to be over-represented in the control tissue. The identified cDNA fragment was found to be a part of the EST clone AF161698  
20 (FIG. 7 B), which encodes the amino acid sequence AAD45360 (FIG. 7 C). This amino acid sequence encodes the Apolipoprotein B mRNA editing protein 2 (APOBEC-2). An overview of the mentioned sequences is depicted in FIG. 7 D.

(APOBEC-2) is highly similar and evolutionarily related to APOBEC-1, which mediates the editing of apolipoprotein (apo) B mRNA (Liao *et al.*). Both proteins are members of C  
25 (cytidine)-->U (uridine) editing enzyme subfamily of the cytidine deaminase supergene family.

APOBEC-2 does not display detectable apoB mRNA editing activity. Like other editing enzymes of the cytidine deaminase superfamily, APOBEC-2 has low, but definite, intrinsic cytidine deaminase activity. APOBEC-2 mRNA and protein are expressed exclusively in  
30 heart and skeletal muscle.

### Yeast two-hybrid interaction

The interaction of AAD45360 (APOBEC-2) was analysed by challenging this bait (against  $4 \times 10^4$  clones). The two-hybrid analysis procedure led to the identification of one interacting partner. This partner was identified by homology search using the first 500 nucleotides sequence of the prey clone. This partner is beta myosin heavy chain (M21665).

The prey cDNA showed 99% homology with beta myosin heavy chain (M21665). Kurabayashi et al., (1988) showed that the beta myosin heavy chain expression is predominantly expressed in the ventricle. Furthermore, the authors show that beta-form MHC mRNA is expressed in adult atrium at a low level but scarcely expressed in fetal atrium. Moreover, mutation of the beta myosin heavy chain have been reported to play a role in heart hypertrophy (Enjuto et al., 2000; Greber-Platzer et al., 2001).

Significant upregulation of 61244 expression in heart tissue of five DCM patients compared to five normal controls indicates that an increased expression of 61244 is associated with dilated cardiomyopathy. Increased expression of 61244 by a factor of 3-4 seems to induce a cardiomyopathic phenotype. Therefore we expect the protein to play a causative role in heart diseases, especially congestive heart failure.

Moreover, the protein is described to be specifically expressed in heart and skeletal muscle. Thus, 61244 may be a novel RNA editing enzyme with natural substrates in these tissues, that plays an important role in RNA modification. This finding supports our idea that 61244 is a specific molecular target for drug development and/or diagnostics.

Downregulation of protein expression by specific inhibitors or antisense constructs seems to be a very promising therapeutic tool to treat heart diseases.

### EXAMPLE 9

Screen for expression profiles in a higher number of patients clearly showed that 65330 is induced upon disease (FIG. 8 E). Transcript levels are significantly upregulated by a factor 2.2 in five DCM patients and 1.8 in two ICM patients compared to five normal controls. The probability of type I error is less than 5% as determined in a Wilcoxon test and t-test.

EST 65330 (FIG. 8 A) was identified by suppression subtractive hybridization comparing transcript levels of heart tissue explanted from normal control KN6 with one from DCM patient h100 (see TABLE 1).

- 5 The identified EST was found to be a part of the EST clone AF249873 (FIG. 8 D), which is itself a part of a 65330contig of assembled EST sequences (FIG. 8 B). The EST clone AF249873 encodes the amino acid sequence AAF63623 (FIG. 8 C).  
AF249873 encodes a novel gene located on human chromosome 4q with specific expression in cardiac and skeletal muscle (Ahmad *et al.*).

10

#### **Yeast two-hybrid interaction**

- 4 x 10<sup>4</sup> clones were challenged against the bait AAF 63623 (SMP). The all two-hybrid analysis procedure led to the identification of one interacting partner:  $\alpha$ -actinin 2 (M86406). This interactor was identified by homology search using the first 500  
15 nucleotides sequence of the prey clone.

#### $\alpha$ -actinin 2

- Homology search with sequences in the database showed 100% identity with  $\alpha$ -actinin 2 (ACTN2) (NM\_001103). The homology starts at nucleotide 1469 of  $\alpha$ -actinin 2.  $\alpha$ -actinin  
20 2 was mapped on chromosome 1q42-q43 and was found to be expressed in skeletal muscle as well as in heart muscle (Beggs *et al.*, 1992).

- Significant upregulation of 65330 expression in heart tissue of five DCM patients and two ICM patients compared to five normal controls indicates that an increased expression of  
25 65330 is associated with dilated cardiomyopathy. According to its interaction with  $\alpha$ -actinin, this protein might play a role in the cytoskeleton of a muscle cell. Therefore we expect the protein to play a causative role in heart diseases, especially in congestive heart failure.

- Moreover, the protein is described to be specifically expressed in heart and skeletal muscle.  
30 This finding supports our idea that 65330 is a specific molecular target for drug

development or diagnostics. Downregulation of protein expression by specific inhibitors or antisense constructs seems to be a very promising therapeutic tool to treat heart diseases.

### EXAMPLE 10

5

EST 66214 (FIG. 9 A) was identified by suppression subtractive hybridization comparing transcript levels of heart tissue explanted from a normal control (KN6) with one from a DCM patient (h100, see TABLE 1). The fragment was found to be over-represented in the DCM tissue.

10 The identified cDNA fragment is a part of the EST clone AF129505; the sequence of the 66214cds is shown in FIG. 9 B.

AF129505 was described to be a novel X-chromosomal human gene (SMPX) encoding the amino acid sequence AAF19343 (9 D) which is a small muscular protein (Patzak *et al.*). The gene consists of five exons and four introns comprising together 52.1 kb and is  
15 preferentially and abundantly expressed in heart and skeletal muscle. The gene maps close to DXS7101 31.9 cM from the short arm telomere of the X-chromosome at Xp22.1. FIG. 9 C shows the amino acid sequence of 66214pep.

Upregulation upon DCM was confirmed for five DCM patients compared to four normal  
20 control hearts by quantitative dot blot analysis (FIG. 9 E). The relative expression level of 66214 is significantly induced by a factor of 4.2 upon disease. The probability of type I error is less than 5% as determined in a Wilcoxon test.

The elevated expression observed for healthy patient h92 may represent individual differences throughout the population.

25

#### Yeast two-hybrid interaction

The  $4 \times 10^4$  clones were analysed for the screen with 66214pep. The two-hybrid analysis procedure led to the identification of 3 different interactors: Daxx (AB015051), Rad6 (U38785), Ubc9 (P50550). These partners were identified by homology search using the  
30 first 500 nucleotides sequence of the pray clone.

Daxx

Search in the data base showed 99% identity with Daxx (AB015051) over the 400 nucleotides. The homology started at nucleotide 1936 of the Daxx sequence. Daxx was mapped on chromosome 6p21.3 (Kiriakidou et al., 1997). The identity found at nucleotide level was confirmed at amino acid level. Daxx was initially found as an interactor of Fas. (Yang et al.1997). Like Fas, it is believed to activate the JNK signal transduction cascade. Therefore, Daxx might play a role in apoptosis regulation.

Ubc9

The prey showed 100% identity with the human Ubc9 sequence. the clone covered the all Ubc9 sequence. Ubc9 is thought to be involved in the ubiquitin-dependent protein degradation system (Wang et al. 1996). A single copy of the hUBC9 gene was found and localised to human chromosome 16p13.3. Interestingly the interaction of Daxx (see above) was already found with the Ubc9 protein (Ryu et al., 2000).

Rad6

Homology search led to the identification of RAD6 (U38785). This result was confirmed by the amino acid analysis. The involvement of RAD6 in the degradation of endogenous inducible cAMP early repressor (ICER) protein in primary cardiomyocytes and myogenic cell lines has been reported (Folco and Koren, 1997 ). Moreover, recent data showed that Ubiquitin-Conjugating Enzymes (rad6) Target Repressors of Cyclic AMP-Induced Transcription for Proteolysis (Pati et al.,1999)

Significant upregulation of 66214 expression in heart tissue of six DCM patients compared to five normal controls indicates that an increased expression of 66214 is associated with dilated cardiomyopathy. Therefore we expect the protein to play a causative role in heart diseases, especially congestive heart failure.

Moreover, the protein is described to be preferentially and abundantly expressed in heart and skeletal muscle. This finding supports our idea that 66214 is a specific molecular target for drug development and/or diagnostics. Downregulation of protein expression by specific

inhibitors or antisense constructs seems to be a very promising therapeutic tool to treat heart diseases.

### EXAMPLE 11

5

66268 and 52474 (FIG. 10 A) were identified by suppression subtractive hybridization comparing transcript levels of heart tissue explanted from normal control KN6 with DCM patient h100, and KN2 with DHZM3 (see TABLE 1), respectively. Both fragments were found to be over-represented in the DCM tissue. Both identified fragments are parts of the  
10 EST clone X83703 (FIG. 10 B), which encodes the amino acid sequence CAA58676 (FIG. 10 C).

CAA58676 has been identified as a novel cytokine-inducible nuclear protein from human endothelial cells (C-193 or CARP, Chu *et al.*). C-193 represents a new member of the  
15 primary response gene family, since its mRNA expression is induced by IL1 $\alpha$ , TNF $\alpha$ , LPS and CHX.

Dot blot hybridizations showed a slight increase in mean expression intensities of DCM patients versus normal controls for both fragments, but the variability from patient to patient was high and the dot blot result statistically was not significant applying a Wilcoxon  
20 or t-test. FIG. 10 E depicts the example of the hybridization with clone 66268.

An overlapping fragment S1MC01-1 was identified to be induced upon DCM by means of differential display (FDD, see 4.). The differential display expression profile independently  
25 confirms upregulation of this gene by a factor of 2.2 upon DCM and ICM and 3.3 upon HCM. The probability of type 1 error for upregulation upon DCM is less than 5% as determined in a t-test.

**Recombinant over expression in primary cardiomyocytes from neonatal rats:**

A CAA58676-YFP fusion protein was over expressed in primary cardiomyocytes from neonatal rats (pCMs). The pCMs were stimulated with Phenylephrine (PE) which leads to flat cells with an extensive parallel sarcomer organization as could be detected in the upper  
5 left and lower right corner of figure 3. The cell over-expressing CAA58676 was detected by the fluorescence signal of the CAA58676-YFP fusion protein. The protein accumulated in little aggregates in the nucleus. In addition, a thin, elongated shape of the cell was detectable, which pointed to the induction of a serial sarcomere organization after over  
10 expression of CAA58676. This observation augmented our opinion, that the over-expression of CAA58676 in the human failing heart has a causative role in disease establishment and progression, because the elongated shape of cardiomyocytes in combination with the serial sarcomere organization is a well known characteristic of diseased cells in the insufficient human heart.

15 Upregulation of 66268 and 52474 expression in heart tissue of DCM, ICM and HCM patients compared to normal controls indicates that an increased expression of 66268 and 52474 is associated with dilated, ischemic and hypertrophic cardiomyopathy. Increased expression of 66268 and 52474 by a factor of 2-3 seems to induce a cardiomyopathic  
20 phenotype. This was strongly supported by our functional analysis in pCMs. A recombinant over expression of a CAA58676-YFP fusion protein led to a serial sarcomere organization which is the main morphological characteristic of diseased cells in the failing human heart. Therefore we expect the protein to play a causative role in cardiomyopathies.

Moreover, the induction by cytokines as well as its mRNA and protein instability elements  
25 indicate an important regulatory function for 66268 and 52474 in signal transduction and control of secondary gene expression. Its ankyrin-like repeats may be involved in protein-protein interactions. These findings support our idea to use 66268 and 52474 as a specific molecular target for drug development and/or diagnostics.

Downregulation of protein expression by specific inhibitors or antisense constructs seems  
30 to be a very promising therapeutic tool to treat heart diseases.

## Claims

1. A method for identifying a subject at risk for a disease of the heart, comprising the step of quantitating the amount of at least one RNA encoding an amino acid sequence selected from the group consisting of:
- 5 (a) the amino acid sequence of SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];
- 10 (b) an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a);
- 15 (c) the amino acid sequence of (a) with at least one conservative amino acid substitution;
- (d) an amino acid sequence that is an isoform of the amino acid sequence of any of (a) to (c);
- 20 (e) the RNA transcribed from the DNA sequence of SEQ ID NO: 10 [NM\_003970], the DNA sequence of SEQ ID NO: 11 [AW755252], the DNA sequence of SEQ ID NO: 12 [EST clone 52706], the DNA sequence of SEQ ID NO: 13 [EST clone 56461], the DNA sequence of SEQ ID NO: 14 [M14780], the DNA sequence of SEQ ID NO: 15 [61166contig], the DNA sequence of SEQ ID NO: 16 [AF161698], the DNA sequence of SEQ ID NO: 17 [65330contig], the DNA sequence of SEQ ID NO: 18 [66214cds] or the DNA sequence AF129505, or the DNA sequence of SEQ ID NO: 19 [X83703] or a degenerate variant thereof; and
- 25 (f) an amino acid that is encoded by a DNA molecule the complementary strand of which hybridizes in 4xSSC, 0.1% SDS at 65°C to the DNA molecule
- 30



encoding the amino acid sequence of (a), (c) or (d),  
in the heart tissue of the subject.

2. The method according to claim 1, wherein the amount of the said RNA is  
5 quantitated using a nucleic acid probe which is a nucleic acid comprising a sequence  
selected from the group consisting of:
- (a) the DNA sequence of the RNA transcribed from the DNA sequence of SEQ  
ID NO: 10 [NM\_003970], the DNA sequence of SEQ ID NO: 11  
[AW755252], the DNA sequence of SEQ ID NO: 12 [EST clone 52706], the  
10 DNA sequence of SEQ ID NO: 13 [EST clone 56461], the DNA sequence of  
SEQ ID NO: 14 [M14780], the DNA sequence of SEQ ID NO: 15  
[61166contig], the DNA sequence of SEQ ID NO: 16 [AF161698], the DNA  
sequence of SEQ ID NO: 17 [65330contig], the DNA sequence of SEQ ID  
NO: 18 [66214cds] or the DNA sequence AF129505, or the DNA sequence  
15 of SEQ ID NO: 19 [X83703] or a degenerate variant thereof;
  - (b) a DNA sequence at least 60%, preferably at least 80%, especially at least  
90%, advantageously at least 99% identical to the DNA sequence of (a);
  - (c) a nucleic acid sequence that encodes the amino acid sequence SEQ ID NO: 1  
[NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the  
20 amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence  
of SEQ ID NO: 4 [AAAS2025], the amino acid sequence of SEQ ID NO: 5  
[61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the  
amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid  
sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence  
25 AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];  
each of said amino acid sequences having at least one conservative amino  
acid substitution;
  - (d) a nucleic acid sequence that encodes an amino acid sequence that is at least  
60%, preferably at least 80%, especially at least 90%, advantageously at  
30 least 99% identical to the amino acid sequence of (c);
  - (e) a nucleic acid sequence that encodes the amino acid sequence of (c) or (d)

with at least one conservative amino acid substitution;

(f) a nucleic acid sequence that hybridizes in 4xSSC, 0.1% SDS at 65°C to the complementary strand of the DNA molecule encoding the amino acid sequence of (c), (d) or (e); and

5 (g) a fragment of at least 15 nucleotides in length of (a) to (f), wherein the nucleic acid is detectably labeled; or

(h) a nucleic acid probe comprising a sequence that specifically hybridizes under physiological conditions to the nucleotide sequence selected from the group consisting of:

10 (i) the DNA sequence of the RNA transcribed from the DNA sequence of SEQ ID NO: 10 [NM\_003970], the DNA sequence of SEQ ID NO: 11 [AW755252], the DNA sequence of SEQ ID NO: 12 [EST clone 52706], the DNA sequence of SEQ ID NO: 13 [EST clone 56461], the DNA sequence of SEQ ID NO: 14 [M14780], the DNA  
15 sequence of SEQ ID NO: 15 [61166contig], the DNA sequence of SEQ ID NO: 16 [AF161698], the DNA sequence of SEQ ID NO: 17 [65330contig], the DNA sequence of SEQ ID NO: 18 [66214cds] or the DNA sequence AF129505, or the DNA sequence of SEQ ID NO: 19 [X83703]

20 (ii) a DNA sequence at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the DNA sequence of (i);

(iii) a nucleic acid sequence that encodes the amino acid sequence SEQ  
25 ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID  
30 NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino sequence of SEQ ID NO: 9 [CAA58676] with at least one

- conservative amino acid substitution;
- (iv) a nucleic acid sequence that encodes an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (iii);
- (v) a nucleic acid sequence that encodes the amino acid sequence of (iii) with at least one conservative amino acid substitution; and
- (vi) a nucleic acid sequence that hybridizes in 2xSSC, 0.1% SDS at 65°C to the DNA molecule encoding the amino acid sequence of (iii), (iv) or (v),
- (vii) a fragment of at least 15 nucleotides in length of (i) to (vi).
3. A method for identifying a subject at risk for a disease of the heart, comprising the step of quantitating the amount of a polypeptide selected from the group consisting of:
- (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];
- (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and
- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution,
- in the heart tissue of the subject.

4. The method according to claim 3, wherein the amount of the said polypeptide is quantitated using an antibody or an antigen-binding portion of said antibody that specifically binds a polypeptide selected from the group consisting of:
- 5 (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961],  
the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid  
sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID  
NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep],  
the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid  
sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ  
10 ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino  
acid sequence of SEQ ID NO: 9 [CAA58676];
- (b) a polypeptide having an amino acid sequence that is at least 60%, preferably  
at least 80%, especially at least 90%, advantageously at least 99% identical  
to the amino acid sequence of (a); and
- 15 (c) a polypeptide having the amino acid sequence of (a) with at least one  
conservative amino acid substitution.
5. The method according to claim 4, wherein said antibody or antibody binding portion  
is or is derived from a human antibody or a humanized antibody.
- 20 6. The method according to claim 4 or claim 5, wherein the antibody, the binding  
portion or derivative thereof is detectably labeled.
7. The method of claim 6, wherein said derivative of said antibody is an scFv  
25 fragment.
8. The method of claim 1 or 2, wherein said RNA is obtained from heart tissue.
9. The method of any one of claims 3 to 7 wherein said polypeptide is quantitated in  
30 heart tissue.

10. The method of any one of claims 1, 2 and 8 further comprising the step of normalizing the amount of RNA against a corresponding RNA from a healthy subject or cells derived from a healthy subject.
- 5 11. The method of any one of claims 3 to 7 and 9 further comprising the step of normalizing the amount of polypeptide against a corresponding polypeptide from a healthy subject or cells derived from a healthy subject.
12. A method for identifying a compound that increases or decreases the level in heart  
10 tissue of a polypeptide selected from the group consisting of:
- (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep],  
15 the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];
- (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical  
20 to the amino acid sequence of (a); and
- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution,
- said method comprising the steps of:
- 25 (1) contacting a DNA encoding said polypeptide under conditions that would permit the translation of said polypeptide with a test compound; and
- (2) detecting an increased or decreased level of the polypeptide relative to the level of translation obtained in the absence of the test compound.
- 30 13. A method for identifying a compound that specifically binds to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 1

[NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; said method comprising the steps of

- (1) providing said polypeptide; and
- (2) identifying a compound that is capable of binding said polypeptide.

14. A monoclonal antibody or derivative thereof that specifically binds to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676].

15. A method for identifying a compound that increases or decreases the level in heart tissue of an mRNA encoding a polypeptide selected from the group consisting of:

- (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];
- (b) a polypeptide having an amino acid sequence that is at least 60%, preferably

at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and

- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution,

said method comprising the steps of

- (1) contacting a DNA giving rise to said mRNA under conditions that would permit transcription of said mRNA with a test compound; and
- (2) detecting an increased/decreased level of the mRNA relative to the level of transcription obtained in the absence of the test compound.

16. A transgenic non-human mammal whose somatic and germ cells comprise at least one gene encoding a functional or disrupted polypeptide selected from the group consisting of:

- (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];

- (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and

- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution,

that said functional or disrupted polypeptide has been modified, said modification being sufficient to decrease or increase the amount of said functional polypeptide expressed in the heart tissue of said transgenic non-human mammal, wherein said transgenic non-human mammal exhibits a disease of the heart.

17. The transgenic non-human mammal according to claim 16, wherein said disrupted or functional gene was introduced into the non-human mammal or an ancestor thereof, at an embryonic stage.
- 5 18. A transgenic non-human mammal according to claim 16 or 17, wherein the modification is inactivation, suppression or activation of said gene(s) or leads to the reduction or enhancement of the synthesis of the corresponding protein(s).
- 10 19. A method for identifying a compound that increases or decreases the expression of a polypeptide in heart tissue, the polypeptide being selected from the group consisting of:
- 15 (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];
- 20 (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and
- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution,
- 25 said method comprising the steps of:
- (1) contacting a transgenic non-human mammal according to any one of claims 14 to 16 with a test compound, and
- (2) detecting an increased or decreased level of expression of said polypeptide relative to the expression in the absence of said test compound.



20. The method according to claim 19, wherein the test compound prevents or ameliorates a disease of the heart in said transgenic non-human mammal.
21. A method for identifying one or a plurality of isogenes of a gene coding for a polypeptide selected from the group consisting of: the polypeptide having the amino acid sequence of SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676]; said method comprising the steps of
- (1) providing nucleic acid coding for said polypeptide or a part thereof; and
  - (2) identifying a second nucleic acid that (i) has a homology of 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% or (ii) hybridizes in 4xSSC, 0.1 SDS at 45°C to the nucleic acid molecule encoding said amino acid sequences.
22. A method for identifying one or a plurality of genes whose expression in heart tissue is modulated by inhibiting, decreasing or increasing the expression of a polypeptide selected from the group consisting of:
- (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];
  - (b) a polypeptide having an amino acid sequence that is at least 60%, preferably

at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and

(c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution,

or of an mRNA encoding said polypeptide,

said modulation being indicative of a disease of the heart, said method comprising the steps of:

- (1) contacting a plurality of heart tissue cells with a compound that inhibits, decreases or increases the expression of said polypeptide under conditions that permit the expression of said polypeptide in the absence of a test compound, and
- (2) comparing a gene expression profile of said heart cell in the presence and in the absence of said compound.

23. A method for identifying one or a plurality of genes whose expression in heart tissue is modulated by the inhibition, decrease or increase of the expression of a polypeptide selected from the group consisting of:

(a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];

(b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and

(c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution,

or of an mRNA encoding said polypeptide, said modulation being indicative of a

disease of the heart, said method comprising the steps of:

- (1) providing expression profiles of
  - (i) a plurality of heart tissue cells from or derived from a heart of a subject  
5 suffering from a disease of the heart; and
  - (ii) a plurality of heart tissue cells from or derived from a subject not suffering from a disease of the heart; and
- (2) comparing the expression profiles (i) and (ii).

10 24. The method of claim 22 further comprising the steps of

- (3) determining at least one gene that is expressed at a lower or higher level in the presence of said compound; and
- (4) identifying a further compound that is capable of raising or lowering the expression level of said at least one gene.

15

25. The method of claim 23 further comprising the steps of

- (3) determining at least one gene that is expressed at a lower or higher level in said heart tissue cells from or derived from a heart of a subject suffering from a disease of the heart; and
- 20 (4) identifying a further compound that is capable of raising or lowering the expression level of said at least one gene.

26. The method of claim 23 further comprising the steps of

- (3) determining at least one gene that is expressed at a higher or lower level in  
25 said heart tissue cells from or derived from a heart of a subject suffering from a disease of the heart; and
- (4) identifying a further compound that is capable of reducing or raising the expression level of said at least one gene.

30 27. A method for identifying a protein or a plurality of proteins in heart tissue whose activity is modulated by a polypeptide having the amino acid sequence selected

from the group consisting of SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];

said method comprising the steps of

- (1) providing said polypeptide; and
- (2) identifying a further protein that is capable of interacting with said polypeptide.

28. The method of any one of claims 12, 13, 15, 19, 20, 22 or 24 to 26, wherein said compound is a small molecule or a peptide derived from an at least partially randomized peptide library.

29. A method of refining a compound identified by the method of any one of claims 12, 13, 15, 19, 20, 22, 24 to 26 or 28; said method comprising the steps of

- (1) identification of the binding sites of the compound and the DNA or mRNA molecule by site-directed mutagenesis or chimeric protein studies;
- (2) molecular modeling of both the binding site of the compound and the binding site of the DNA or mRNA molecule; and
- (3) modification of the compound to improve its binding specificity for the DNA or mRNA.

30. The method of any one of claims 12, 13, 15, 19, 20, 22, 24 to 26, 28 or 29, wherein said compound is further refined by peptidomimetics.

31. A method of modifying a compound identified or refined by any one of claims 12, 13, 15, 19, 20, 22, 24 to 26, 28 to 30 as a lead compound to achieve

- (i) modified site of action, spectrum of activity, organ specificity, and/or  
(ii) improved potency, and/or  
(iii) decreased toxicity (improved therapeutic index), and/or  
(iv) decreased side effects, and/or  
5 (v) modified onset of therapeutic action, duration of effect, and/or  
(vi) modified pharmacokinetic parameters (resorption, distribution, metabolism and excretion), and/or  
(vii) modified physico-chemical parameters (solubility, hygroscopicity, color, taste, odor, stability, state), and/or  
10 (viii) improved general specificity, organ/tissue specificity, and/or  
(ix) optimized application form and route

by

- (i) esterification of carboxyl groups, or  
15 (ii) esterification of hydroxyl groups with carbon acids, or  
(iii) esterification of hydroxyl groups to, e.g. phosphates, pyrophosphates or sulfates or hemi succinates, or  
(iv) formation of pharmaceutically acceptable salts, or  
(v) formation of pharmaceutically acceptable complexes, or  
20 (vi) synthesis of pharmacologically active polymers, or  
(vii) introduction of hydrophylic moieties, or  
(viii) introduction/exchange of substituents on aromates or side chains, change of substituent pattern, or  
(ix) modification by introduction of isosteric or bioisosteric moieties, or  
25 (x) synthesis of homologous compounds, or  
(xi) introduction of branched side chains, or  
(xii) conversion of alkyl substituents to cyclic analogues, or  
(xiii) derivatisation of hydroxyl group to ketals, acetals, or  
(xiv) N-acetylation to amides, phenylcarbamates, or  
30 (xv) synthesis of Mannich bases, imines, or  
(xvi) transformation of ketones or aldehydes to Schiff's bases, oximes, acetals,

ketales, enolesters, oxazolidines, thiozolidines  
or combinations thereof.

32. A method for inducing a disease of the heart in a non-human mammal, comprising  
5 the step of contacting the heart tissue of said mammal with a compound that  
inhibits, decreases or increases the expression of a polypeptide selected from the  
group consisting of:
- (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961],  
the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid  
10 sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID  
NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep],  
the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid  
sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ  
ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino  
15 acid sequence of SEQ ID NO: 9 [CAA58676];
  - (b) a polypeptide having an amino acid sequence that is at least 60%, preferably  
at least 80%, especially at least 90%, advantageously at least 99% identical  
to the amino acid sequence of (a); and
  - (c) a polypeptide having the amino acid sequence of (a) with at least one  
20 conservative amino acid substitution.
33. The method according to claim 32, wherein said compound that inhibits, decreases  
or increases is a small molecule, an antibody or an aptamer that specifically binds  
said polypeptide.
- 25 34. A method of producing a pharmaceutical composition comprising formulating the  
compound identified, refined or modified by the method of any of the preceding  
claims with a pharmaceutically active carrier or diluent.
- 30 35. A method for preventing or treating a disease of the heart in a subject in need of  
such treatment, comprising the step of increasing or decreasing the level of a

polypeptide in the heart tissue of a subject, said polypeptide being selected from the group consisting of:

- 5 (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];
- 10 (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and
- 15 (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution.

36. A method of preventing or treating a disease of the heart in a subject in need of such treatment comprising the step of increasing or decreasing the level of mRNA encoding a polypeptide in the heart tissue of a subject, said polypeptide being selected from the group consisting of:

- 20 (a) the polypeptide having amino acid sequence SEQ ID NO: 1 [NP\_003961], the amino acid sequence of SEQ ID NO: 2 [41441pep], the amino acid sequence of SEQ ID NO: 3 [56461pep], the amino acid sequence of SEQ ID NO: 4 [AAA52025], the amino acid sequence of SEQ ID NO: 5 [61166pep], the amino acid sequence of SEQ ID NO: 6 [AAD45360], the amino acid sequence of SEQ ID NO: 7 [AAF63623], the amino acid sequence of SEQ ID NO: 8 [66214pep] or the amino acid sequence AAF19343, or the amino acid sequence of SEQ ID NO: 9 [CAA58676];
- 25 (b) a polypeptide having an amino acid sequence that is at least 60%, preferably at least 80%, especially at least 90%, advantageously at least 99% identical to the amino acid sequence of (a); and
- 30

- (c) a polypeptide having the amino acid sequence of (a) with at least one conservative amino acid substitution.

37. The method of claims 35 or 36, wherein such increase or decrease is effected by administering the pharmaceutical composition obtained by the method of claim 30.
38. The method of claim 35 or 36, wherein such an increase or decrease is effected by introducing the nucleic acid sequence recited in claim 2 into the germ line or into somatic cells of a subject in need thereof.
39. The method of any of the preceding claims, wherein said disease of the heart is congestive heart failure, dilative cardiomyopathy, hypertrophic cardiomyopathy, ischemic cardiomyopathy, specific heart muscle disease, rhythm and conduction disorders, syncope and sudden death, coronary heart disease, systemic arterial hypertension, pulmonary hypertension and pulmonary heart disease, valvular heart disease, congenital heart disease, pericardial disease or endocarditis.
40. A method for identifying subjects at risk for heart diseases, especially congestive heart failure, comprising the step of detecting an increased or decreased level of MYOM2, the LIM domain, the muscle isoform of creatine kinase, YAP65, APOBEC-2, SMPX or C-193 (CARP) in the heart tissue of a subject.
41. A method for preventing or treating heart diseases, especially congestive heart failure in a subject, said method comprising the step of contacting the heart tissue of said subject with a compound that decreases or increases the expression of MYOM2, the LIM domain, the muscle isoform of creatine kinase, YAP65, APOBEC-2, SMPX or C-193 (CARP).
42. A method for identifying subjects at risk for heart diseases, especially congestive heart failure, comprising the step of detecting decreased creatine kinase activity in the tissue of a subject especially in a muscle tissue or from blood or serum.



- 5
43. A method for identifying a subject at risk for heart diseases, especially congestive heart failure, said method comprising detecting increased levels of creatine phosphate in a subject, especially in the blood or serum of a subject.
44. A method for preventing or treating heart diseases, especially congestive heart failure in a subject, said method comprising the step of increasing the transfer of phosphoryl groups from creatine phosphate to ADP in the heart tissue of a subject.
- 10 45. The method according to claim 44, wherein the activity of creatine kinase is increased in said heart tissue.
46. A method for identifying a compound for preventing or treating heart diseases, especially congestive heart failure, said method comprising the steps of:
- 15
- (a) contacting creatine kinase with a substrate for creatine kinase and a test compound, and
  - (b) determining whether the transfer of phosphoryl groups from the substrate is increased in the presence of the test compound.
- 20
47. Use of a compound of one of the claims 12, 13, 15, 19, 20, 24 to 26, 28, 41, 46, a refined or modified compound of one of the claims 29, 30 or 31, or a monoclonal antibody of the claim 14 for the manufacture of a pharmaceutical composition for the prophylaxis or treatment of heart diseases, especially congestive heart failure.
- 25

Length: 197 nt  
>40399  
1 ACAGACGAAA TGAAAGTGAA CTGGTGTAC AAAGATGCTC AGACCTCATC CAGTGAGCAT  
61 ATGAGAAATCC GGGGAGTGA AGAGATGGCT TGGCTGCACA TATGTGAGCC GACTGACAAG  
121 GATAAAGGAA AATACACTTT TGAGATTTTC GATGGCTAAT ACAACCATCA ACGCTCCCTT  
181 GACCTGTCCG GACACTG

FIG. 1 A

## FIG 1 B/1

Length: 4939 nt  
 >NM\_003970

1	ttctctctctcc	tccttgcaat	tttcctttct	gtctgggagc	acgccaagat	gtcccttgtg
61	actgtccccc	tctaccagaa	gagacatagg	cacttcgacc	agtcctaccg	taatatccaa
121	acacgggtacc	tgctggacga	atatcggtca	aaaaagcgag	cttccaccca	ggcatctctcc
181	cagaagtcct	tgagtcagcg	gtcgtcttca	cagagagcct	ccagccagac	gtccctggga
241	ggaaccatct	gcagggtctg	tgcgaagcga	gtgagcacgc	aggaagatga	ggagcaggag
301	aacagaagca	ggtaccagtc	cctgggtggcc	gcctatggtg	aggccaagcg	acacggcttc
361	ctcagcgagc	tggcccactt	ggaggaggat	gtccacctgg	cacgctccca	ggcccgcgac
421	aagctggaca	aatacgccat	tcagcagatg	atggaggaca	agctggcctg	ggagagacac
481	acatttgaag	agcggataag	cagggtctct	gagatcctgg	tgcggctgcg	atcccacacc
541	gtctgggaga	ggtgtctgt	gaaactctgc	ttcacctgtc	aaggatttcc	cacgcccgtg
601	gtgcagtggc	acaaagatgg	cagtctgatt	tgccaggcgg	ctgaaccggg	aaagtacagg
661	attgagagca	actatggcgt	acacacactg	gagatcaaca	gggcagactt	tgacgacact
721	gcgacatact	cagcagtggc	caccaatgcc	cacggacaag	tgtccacca	cgcggcgggtg
781	gtgggtgagaa	ggttccgggg	agacgaggaa	ccattccgtt	cggaggact	cccgattgga
841	ttgcccctgt	catcgatgat	tccgtacacg	cacttcgacg	tccagttttt	ggagaaagt
901	gggggtcacct	tcaggaggga	aggcgagacg	gtcactctca	agtgcaccat	gctggtgacg
961	ccgggacctga	agcgggtgca	gccgcgcgcc	gagtgggtacc	gcgatgactt	gctggtgaaa
1021	gagtcctcaagt	ggacgaagat	gttcttttga	gaaaggccagg	cctccctgtc	cttcagccac
1081	ctgcacaaagg	acgacgaggg	cctgtacacc	ctgcgcacatcg	tgtctcgggg	cggcgtcacg
1141	gaccacagcg	ccttcctgtt	tgtcagagat	gctgacccgc	tggtcacagg	ggcccccggt

FIG. 1 R/2

```

1201 gcacccatgg acttgcaagt ccacgacgcc aaccgggact acgtcatcgt gacctggaag
1261 ccgcccaca ccaccactga gagccccgtc atgggctatt ttgtggaccg atgtgaagta
1321 ggaacgaata attgggtgca gtgcaatgat gcaccggtga aaatctgcaa ataccgggtc
1381 acagggcctt ttgaagggaag gtcttacata ttccgagtga gggcagtgaa cagtgcgggc
1441 atcagccgac cctccagggt ctctgatgcg gtggctgcac ttgacccctt ggacctcaga
1501 aggttacaag ccgttcattt ggagggagag aaggagattg ccatttatca ggatgacctt
1561 gaaggtgacg ccaggttcc ccagggttcc accggtgtgc acgcttcga gatcagcaga
1621 aactatgtcg tcctcagctg ggagccaccc actcccgtg gcaaggaccg gctcatgtac
1681 ttcattgaga agtcggtggt ggggagcggc acgtggcaga gagtcaacgc ccagacggct
1741 gtgagatccc cgagatatgc cgtgtttgac ctcatggaa ggaagtctta tgtgttccga
1801 gtgctgtcag caaacgggca tggcctgagc gaaccttcgg agataaacgc cccattcag
1861 gccaggatg tgaccgttgt cccttcgtct ccgggtcggg ttcttgcttc ccgaaacacc
1921 aagacgtcgg tggtagtgca gtgggaccga cctaagcatg aggaggacct gctgggctac
1981 tacgtggact gctgtgtggc cggaaaccaac ctctgggagc cctgcaacca caagcccatc
2041 ggatacaaca ggttcgtggt gcacggctta accacgggag agcagttacat ctccgagtc
2101 aaggcgggtca atgctgtggg gatgagtga aattcccagg aatcagacgt cataaaagtg
2161 caggccgcac tcaccgtccc gtcccattct tatgggatta cgctcctcaa ctgtgacggc
2221 cactccatga ccctcggctg gaagggtccc aaattcagtg gtggctcgc catcctgggc
2281 tactacctgg acaagcgtga agttcaccat aaaaactggc acgaggtcaa ttctcacccc
2341 agcaaaccca caatcctaac ggtggacggc ttgacgggaag gctcactcta cgagttcaaa
2401 atcgccgccg tcaacctggc cggcatcggg gagccctcag atcccagtga gcacttcaag
2461 tgtgaggcct ggaccatgcc ggagcccggt cctgcctacg acttgacgtt ctgtgaggtc

```

FIG. 1 B/3

2521 agggacacgt ccttggtcat gctgtggaag gccctgtgt actccggcag cagccctgtt  
2581 tctggatat tctggactt cagggaggag gatgctggag agtggatcac tgtcgatcac  
2641 acgacaacag ccagccgtta tttaagggtc tctgacctgc agcaaggtaa gacctatgtc  
2701 ttcaggggtc gggcagtc aa ggcagagacc aggcaccaag gaaatcagtg ctcagacac gtcggagcct  
2761 gtgctggtag aggcagagacc tgggcttcga agatgatgag aggttttaaa atgacagacg cgtctcagtt cacctggtgt  
2821 aacatctatc tgggcttcga aggcagagacc aggcagagacc aggttttaaa atgacagacg cgtctcagtt cacctggtgt  
2881 aaatcctacg aggcagagacc aggcagagacc aggcagagacc aggttttaaa atgacagacg cgtctcagtt cacctggtgt  
2941 tccaagctgt acttaagaa tccggataag gaggatttag gacttactc tcgaaaccgt gggggtcac  
3001 agtgatatcag acggagtgtc tccagttttt gttctggacc cagaagagct cgagcgtttg  
3061 atggcatgga gcaatgaaat aaagaacccc acaattcctc tgaatcggg attagcttat  
3121 gagatttttg ataaggggag ggctccagg tggagcactt atcaccagat  
3181 gccagctacc gatttattat taatgacaga gaagtctctg acagcgagat acacagaaat  
3241 aaatgtgaca aagctactgg cattattgag atggtgatgg atcgatttag tattgaaaat  
3301 gaggggacct acactgtgca gattcatgat gggaaagcca aaagtcagtc ttctctagtt  
3361 ctatttggag atgcattcaa gactgtgctg gaagaggctg agtttcaaag gaaagaattt  
3421 ctcaaggaaac aaggccctca ttttgctgag tacttgcaat accaagaaag gggatgtcac ggaagaatgt  
3481 gaagtctgac ttgtttgcaa ggttgcaaac accaagaaag aaaccgtttt caaatggctc  
3541 aaggatgatg ctctgtatga aacggagaca ctgacctaac tggagagggg aatctgtgag  
3601 ctccctcatcc caaagtgtgc aaagaaggac cacggtgaat acaaggcaac cttgaaagat  
3661 gacagaggcc aagatgtgtc catccttgaa atagctggca aagtgtatga tgatatgat  
3721 ttggcaatga gtagagtctg tgggaaatct gcttcgccac tgaagggtact ctgcaccca  
3781 gaaggaatac gacttcagtg ttatcatgaag tattttacag acgaaatgaa agtgaaactgg

FIG. 1 B/4

```

3841  tgtcacaaag  atgctaagat  ctcatccagt  gagcataatga  gaatcggggg  gagtgaagag
3901  atggcttggc  tgcagatatg  tgagccgact  gagaaggata  aagggaaaata  cacttttgag
3961  attttcgatg  gcaaagacaa  ccatcaacgc  tcccttgacc  tgtccggaca  agcttttgat
4021  gaagcatttg  cagaattcca  gcaattcaaa  gctgctgctt  ttgcagagaa  gaatcgtggc
4081  aggttgatcg  gcggcttgcc  tgacgtgggtg  accatcatgg  aagggaagac  cttgaatctg
4141  acctgcacgg  tgtttggaaa  ccctgacccc  gaagtgattt  ggttcaagaa  cgaccaggac
4201  atccagctca  gcgagcactt  ctcggtgaag  gtggagcagg  ccaagtacgt  cagcatgacc
4261  atcaaaaggc  tgacctccga  ggactcgggc  aagtacagca  tcaacatcaa  gaataagtat
4321  ggcggggaga  agatcgacgt  gacgtgagc  gtgtacaaac  acggggagaa  gatcccggac
4381  atggccccc  ccagcaagc  caagcccaag  ctcatccccg  cgtctgcctc  agcggcaggc
4441  cagtgaaggc  gttttccctag  cctggagatg  ggaaaaatatg  ctgggcagag  acaggaaatgc
4501  tgtgtgcttg  ttccaaatga  gcagctggca  tccgagtgggt  gtccctgtgtg  ggctgatatg
4561  tgatcacaca  ttgtgctttt  gattttttgca  ttgggtgatg  aatattttat  acccgtctaa
4621  gggagaaagc  taatgttttc  cacaagactg  aacaacgtgt  atttacacga  gggtagacgg
4681  cagatgcctg  acagagagtg  ggttggcaga  caacacacta  gcattttcac  ggggtgtggc
4741  acatgggtgt  ggcacctgga  cgtgtgcagc  atgtggcgggt  ctctgtgtga  agccaccgtg
4801  ctctctttg  gggggccgcg  agatctagca  tctctgaaat  cctggctgtc  gaggctttga
4861  agcatgtgtt  acctggttaa  gcttgtttc  tcttgcttta  ggcaataaaa  agtttaaaaa
4921  tcaaaaaaaa  aaaaaaaaaa

```

Length: 1465 amino acids


>NP\_003961

```

MSLVTVPFYQ KRHRHFDQSY RNIQTRYLLD EYASKKRAST QASSQKSLSQ 51
RSSSQRASSQ TSLGGTICRV CAKRVSTQED EEQENRSRYQ SLVAAYGEAK 101
RHGFLSELAH LEEDVHLARS QARDKLDKYA IQQMEDKLA WERHTFEERI 151
SRAPEILVRL RSHTVWERMS VKLCFTVQGF PTPVVQWYKD GSLICQAAEP 201
GKYRIESNYG VHTLEINRAD FDDTATYS AV ATNAHGQVST NAAVVVRRFR 251
GDEEPFRSVG LPIGLPLSSM IPYTHFDVQF LEKFGVTFRR EGETVTCLKCT 301
MLVTPDLKRV QPRAEWYRDD LLLKESKWK MFFGEGQASL SFSHLHKDDE 351
GLYTLRIVSR GGVTDHSAFL FVRDADPLVT GAPGAPMDLQ CHDANRDYVI 401
VTWKPPNTTT ESPVMGYFVD RCEVGTTNNWV QCNDAPVKIC KYPVTGLFEG 451
RSYIFRVRAV NSAGISRPSR VSDAVAALDP LDLRRLQAVH LEKEKEIAIY 501
QDDLEGDAQV PGPPTGVHAS EISRNYVVL S WEPPTPRGKD PLMYFIEKSV 551
VGSGTWQRVN AQTAVRSPRY AVFDLMEGKS YVFRVLSANR HGLSEPSEIT 601
SPIQAQDVT VPSAPGRVLA SRNTKTSVVV QWDRPKHEED LLGYVDDCCV 651
AGTNLWPCN HKPIGYNRFV VHGLTTGEQY IFRVKAVNAV GMSSENSQESD 701
VIKVQAALTV PSHPYGITLL NCDGHSMTLG WKVPKFSGGS PILGYLDKR 751
EVHHKNWHEV NSSPSKPTIL TVDGLTEGSL YEFKIAAVNL AGIGEPSDPS 801
EHFKCEAWTM PEPGPAYDLT FCEVRDTSLV MLWKAPVYSG SSPVSGYFVD 851
FREEDAGEWI TVDQTTTASR YLKVSDLQQG KTYVFRVRAV NANGVGKPSD 901
TSEPVLVEAR PGTKEISAGV DEQGNIIYLG DCQEMTDASQ FTWCKSYEEI 951
SDDERFKIET VGDHSLKLYLK NPDKEDLGT SVSVSDTDGV SSSFVLDPEE 1001
LERLMALSNE IKNPTIPLKS ELAYEIFDKG RVRFWLQAEH LSPDASYRFI 1051
INDREVSDSE IHRKCDKAT GIEMVMDFR SIENEGTYTV QIHDGKAKSQ 1101
SSLVLIGDAF KTVLEEAFFQ RKEFLRKQGP HFAEYLHWDV TEECEVRLVC 1151
KVANTKKETV FKWLKDDALY ETETLPNLER GICELLIPKL SKKDHGEYKA 1201
TLKDDRGQDV SILEIAGKVY DDMILAMSRV CGKSASPLKV LCTPEGIRLQ 1251
CFMKYFTDEM KVNWCHKDAK ISSSEHMRIG GSEEMAWLQI CEPTEKDKGK 1301
YTFEIFDGKD NHQSLDLG QAFDEAFAEF QQFKAAAF AE KNRGRLIGGL 1351
PDVVTIMEGK TLNLTCTVFG NPDPEVIWFK NDQDIQLSEH FSVKVEQAKY 1401
VSMTIKGVTS EDSGKYSINI KNKYGGEKID VTVSVYKHGE KIPDMAPPQQ 1451
AKPKLIPASA SAAGQ

```

FIG. 1 C



40399 (197 nt)

NM\_003970 (4939 nt)  
identical to X69089

NP\_003961 (1465 aa)  
identical to CAA48832

FIG. 1 D






FIG. 1 E

Length: 403 nt  
>41441  
1 AAGAAGAAGA GATGTGCAAG GATAGGCCGA GTGAAGCTGA AGACACAAAG AGTACAGGAA  
61 AAGTGCTATG GATCTTAATG ACAACAATAA TGTGATTGTG CAGAGTGCTG AAAAGGAGAA  
121 AAATGAAAAA ACTAACCAAA CTAATGGTGC AGAAGTTTAA CAGGTTACTA ACACTGATGA  
181 TGAGATGTGC CAGAAAATCA TAAAGAAAAT TTGAATAAGA ATAATAATA CAATTATGTA  
241 GCAGTCTCAT ATCTGAATAA TTGCAGGCAG AAGACATCTA TTTTAGAATT TCTTGATCTA  
301 TTACCCCTGT CGAGTGAAGC AAATGACACT GCAAATGAAT ATGAAATTGA GAAGTTAGAA  
361 AATACATCTA GAATCTCAGA GTTACTTGGT AGATTTGAAT CTG

FIG. 2 A

Length: 2379 nt  
>AW755252

```

1  ccaggatct  gctctgaaac  caggtctcta  agtgaacatt  tctcaggcat  ggatgcattt
61 gagagtcaaa  ttgttgagtc  gaagatgaaa  acctcttcac  cacatagctc  agaagctggc
121 aaatctggct  gtgacttcaa  gcatgcccc  ccaacctatg  aggatgtcat  tgctggacat
181 attttagata  tctctgattc  acctaaagaa  gtaagaaaaa  attttcaaaa  gacgtggcaa
241 gagagtggaa  gagtttttaa  aggcctggga  tatgcaaccg  cagatgcttc  tgcaacatga
301 gatgagaacc  accttccaag  aggaatctgc  atttataagt  gaagctgctg  ctccaagaca
361 aggaatatg  tatactttgt  caaaagacag  tttatccaat  ggagtgccta  gtggcagaca
421 agcagaattt  tcataagtcc  tgcttccgat  gccaccattg  caacagtaaa  ctaagtttgg
481 gaaattatgc  atcacttcac  ggacaaaatat  actgtaaac  tcaacttttca
541 aatccaaagg  aaatttatgat  gaaggttttg  gacataagca  gcataaagat  agatgggaact
601 gcaaaaaacca  aagcagatca  gtggacttta  ttcctaataga  agaaccaaat  atgtgtaaaa
661 atattgcaga  aaacaccctt  gtacctggag  atcgtaataga  acattttagat  gctggtaaca
721 gtgaagggca  aaggaatgat  ttgagaaaaa  taggggaaag  gggaaaaatta  aaagtcattt
781 ggcctccttc  caaggagatc  cctaagaaaa  ccttaccctt  tgaggaaagag  ctcaaaatga
841 gtaaacctaa  gtggccacct  gaaatgacaa  ccctgctatc  ccctgaattt  aaaagtgaat
901 ctctgctaga  agatgtttaga  actccagaaa  ataaaggaca  aagacaagat  cacttttccat
961 ttttgcagcc  ttatctacag  tccaccccatg  tttgtcagaa  agaggatgtt  ataggaatca
1021 aagaaatgaa  aatgcctgaa  ggaagaaaaag  atgaaaaagaa  ggaaggaaagg  aagaatgtgc
1081 aagataggcc  gagtgaagct  gaagacacaa  agagtaacag  gaaaaagtgc  atggatctta

```

FIG. 2 B/1

```

1141 atgacaacaa taatgtgatt gtgcagagtg ctgaaaagga gaaaaatgaa aaaactaacc
1201 aaactaatgg tgcagaagtt ttacaggtta ctaacactga tgatgagatg atgccagaaa
1261 atcataaaga aaatttgaat aagaataata ataacaatta ttagcagtc tcatatctga
1321 ataattgcag gcagaagaca tctattttag aatttcttga tctattacc ttgtcgagtg
1381 aagcaaatga cactgcaaat gaatatgaaa ttgagaagtt agaaaaataca tctagaatct
1441 cagagttact tggatatatt gaactgaaa agacttattc gaggaatgta ctagcaatgg
1501 ctctgaagaa acagactgac agagcagctg ctggcagtc ctggcagcct gctccaaaaac
1561 caagcctcag cagaggcctt atggtaaaagg ggggaagtcc aatcatctct cctgatacaa
1621 atctcttaaa cattaaagga agccattcaa agagcaaaaa ttacacttt ttcttttcta
1681 acaccgtgaa aatcactgca tttccaaga aaaatgagaa cattttcaat tgtgatttaa
1741 tagattctgt agatcaaat aaaaatatgc catgcttgggaa tttaagggaa ttgggaaagg
1801 atgttaaac ttggcatgtt gaaacaacag aagctgcccg caataatgaa aacacaggtt
1861 ttgatgctct gagccatgaa tgtacagcta agcctttgtt tccagagtg gaggtgcagt
1921 cagaacaaact cacggtgga gagcagatta aaagaaacag gtgctacagt gacactgagt
1981 aaaatatcta tggccactga cagtccacac ttagggcactg agagatatg atgttctgaa
2041 ataagatttt atgaatttgg ataccctttt gaggaacttg atgtaaacat ggtgttcaga
2101 aatctcgtgt ctatctcaat gggatatattc ttgtattaca ccttgtcatt tttttcaciaa
2161 ttattttaca tctacttttg tttgaaactgg aatgaagaga tgaaacacta tggatatgtt
2221 tccattcaa atggcacttt agcatattgt tctgttttcc tgtaaacat catgggtgtg
2281 atttttatag tgctgctgct tgtcacaatt attataactt ctctgtaatt tcctctgaaa
2341 taaaattgaa tcacctgagg tgcaaaccaa aaaaaaaa

```

FIG. 2 B/2

Length: 547 amino acids  
>41441pep

1 VKLLQLQDKEI CILCQKTVYP MECLVADKQN FHKSCFRCHH CNSKLSLGNY ASLHGQIYCK  
61 PHFKQLFKSK GNYDEGFGHK QHKDRWNCKN QSRSVDFIPN EEPNMCKNIA ENTLPVPGDRN  
121 EHL DAGNSEG QRNDLRKLGE RGKLKVIWPP SKEIPKKTLP FEEELKMSKP KWPPEMTTLL  
181 SPEFKSESLL EDVRTPENKG QRQDHFPPFLO PYLQSTHVCQ KEDVIGIKEM KMPEGKDEK  
241 KEGRKNVQDR PSEAEDTKSN RKSAMDNDN NNVIQSAEK EKNEKTNQTN GAFLVQVTNT  
301 DDEMPENHK ENLNKNNNN YVAVSYLNNC RQKTSILEFL DLLPLSSEAN DTANEYEIEK  
361 LENTSRISEL LGIFESEKTY SRNVLAMALK KQTDRAAAGS PVQPAPKPSL SRGLMVKGGS  
421 SIISPDTNLL NIKGSHSKSK NLHFFFSNTV KITAFSKKNE NIFNCDLIDS VDQIKNMPCL  
481 DLREFGKDVK PWHVETTEAA RNNENTGFDA LSHECTAKPL FPRVEVQSEQ LTVEEQIKRN  
541 RCYSDTE

FIG. 2 C





FIG. 2 D




heart tissue sample

FIG. 2 E

Length: 125 nt  
>52706  
1 ACAGCTTACA GAACGTGGG CCAATAAAC CTCCTTCTT TATAAATTAC CCAGCTTCAG  
61 ATATTCCTTT ATAGCGACAC AAATGGACTA AGGTGTCAAG ATCATTTGAT AGAGAAAGGC  
121 ATTGT

FIG. 3 A





heart tissue sample

FIG. 3 B

Length: 508 nt  
>56461

```
1  GGCTGCATAG TCTTGGCGGA GGTGACCAAA GCCGCGTAAT GTCCGTAGTT CGCTCATCCG
61  TCCATGCCAG ATGGATTGTG GGAAGGTGA TTGGGACAAA AATGCAAAAG ACTGCTAAAG
121 TGAGAGTGAC CAGGCTTGTT CTGGATCCCT ATTTATTAAA GTATTTTAAT AAGCGGAAAA
181 CCTACTTTGC TCACGATGCC CTTCAGCAGT GCACAGTTGG GGATAATTGT CTTCTCAGAG
241 CTTTACCTGT TCCACGAGCA AAGCATGTGA AACATGAACT GGCTGAGATC GTTTTCAAAG
301 TTGGAAAAGT CATAGATCCA GTGACAGGAA AGCCCTGTGC TGGAACTACC TACCTGGAGA
361 GTCCGTTGAG TCGGAAACCA CCCAGCTAAG CAAAATCTG GAAGAACTCA ATATCTCTC
421 AGCACAGTGA AGCGGGAGTG AAGAAGGATC TAAAGGGAAA AACTGACATG TTTATGTTAT
481 GGAAAAAGAA ATTTTCTTAA GTTTCATC
```

FIG. 4 A

Length: 600 nt  
>AF077035

```
1 agtcttggcg gagtgacca aagcacgta atgtccgtag ttcgctcatc cgtccatgcc
61 agatggattg tggggaagg gattgggaca aaaatgcaaa agactgctaa agtgagagtg
121 accaggcttg ttctggatcc ctatttatta aagtatttta ataaaggga aacctacttt
181 gctcacgatg cccttcagca gtgcacagtt ggggatatatg tgcttctcag agctttacct
241 gtccacgag caaagcatgt gaaacatgaa ctggctgaga tcgttttcaa agttggaaaa
301 gtcatagatc cagtgcagag aaagccctgt gctggaacta cctaccctgga ggtccgttg
361 agttcggaaa ccaccagct aagcaaaaat ctggaagaac tcaatatctc tcagcacag
421 tgaagcggga gtggaagaag ggtctaaagg gaaaaactga catgtttatg ttatggaaaa
481 agaaaatttt ctaagtttca tcacaaactg gtccagttt ctctgtggtg ttatgaaat
541 agctaaaaagc aaatgaagta aagggcatac tatggttttt cacaaaaaaa aaaaaaaaaa
```

FIG. 4 B

Length: 142 amino acids  
>56461pep

1 LHSLGGDQS HVMVVRSSV HARWIVGKVI GTKMQKTAKV RVTRLVLDPY LLKYFNKRKT  
61 YFAHDALQQC TVGDI~~V~~LLRA LPVPRAKHVK HELAEIVFKV GKVIDPVTGK PCAGTTYLES  
121 PLSSETTQLS KNLEELNISS AQ

FIG. 4 C





FIG. 4 D

56461 (508 nt)

AF077035 (600 nt)

AAD27768 (130 aa)

56461 pep



heart tissue sample

FIG. 4 E

Length: 383 nt  
>61105  
1 ACCCCTCTTC TGCAGACGCA GCGGGTGAG GATCTCCTCG AACTTGGGGT GCTTGCTCAG  
61 GTGCGCCAGC TTCACATGCA CGCCTCCACG CAGCCAGTG CCCAGGTGG ATGGGCAGGT  
121 GAGCACGTAG CCCAGGTGCT GGTTCACAT GAAGGGGTGG CCAGCTTTCT TAAAGATCTC  
181 CTCATCTTC TGCAGCCCTA CGCAGAAGCG GCGGAAACC TCCTTCATGT TGCCCCCCTT  
241 CTCCATGGAG ATGACCCGGA GGTGATCCTC CTCGTCACC CACACCAGGA AGCTCTTGT  
301 GTCATTGTGG CAGATGCCAC GGGCGTCGG CCATGCCCTG AGGCCAGCAG  
361 CAGCGGGGA CACGGGCTTT GTC

FIG. 5 A

Length: 1562 nt  
>M14780

```

1  gtgggtcagc  atgtcacctc  caggatacag  acagccccc  ttacgccag  ccagccagg
61  tctccttaca  ccgccaccat  gccattcggg  aacacccaca  acaagttcaa  gctgaattac
121 aagcctgagg  aggagtaccc  cgacctcagc  aacataaaca  accacatggc  caaggtactg
181 acccttgaac  tctacaagaa  gctgcgggac  aaggagatcc  catctggctt  cactgtagac
241 gatgtcatcc  agacaggagt  ggacaaccca  ggtcacccct  tcatcatgac  cgtgggctgc
301 gtggctggtg  atgaggagtc  ctacgaagtt  ttcaaggaa  tctttgacct  catcatctcg
361 gatcgccacg  ggggctacaa  acccactgac  aagcacaaga  ctgacctcaa  ccatgaaaac
421 ctcaagggtg  gagacgacct  ggaccccaac  tacgtgctca  gcagcccggt  ccgcactggc
481 cgcagcatca  agggctacac  gttgccccca  cactgctccc  gtggcgagcg  ccgggcggtg
541 gagaagctct  ctgtggaagc  tctcaacagc  ctgacgggcg  agttcaaagg  gaagtactac
601 cctctgaaga  gcatgacgga  gaaggagcag  cagcagctca  tcgatgacca  cttccagttc
661 gacaagcccg  tgtccccgct  gctgctggcc  tcaggcatgg  ccgcccactg  gcccgacgcc
721 cctggcatct  ggcacaaatga  caacaagagc  ttccctggtgt  ggggtgaacga  ggaggatcac
781 ctccgggtca  tctccatgga  gaaggggggc  aacatgaagg  aggttttccg  ccgcttctgc
841 gtagggctgc  agaagattga  ggagatcttt  aagaaagctg  gccacccctt  catgtggaac
901 cagcacctgg  gctacgtgct  cacctgcccc  tccaacctgg  gcactgggct  gcgtggaggc
961 gtgcatgtga  agctggcgca  cctgagcaag  caccacaagt  tcgaggagat  cctcacccgc
1021 ctgcgtctgc  agaagagggg  tacaggtgcg  gtggacacag  ctgccgtggg  ctcagtattt
1081 gacgtgtcca  acgctgacgc  gctgggctcg  tccgaagtag  aacaggtgca  gctggtggtg

```

FIG. 5 B/1



1141 gatggtgtga agctcatggt ggaaatggag aagaagttag agaaaggcca gtccatcgac  
1201 gacatgatcc ccgccagaa gtaggcgct gccacctgc caccgactgc tggaaaccca  
1261 gccagtggga gggcctggcc caccagagtc ctgctccctc actcctcgcc ccgccccctg  
1321 tcccagagtc cacctggggg ctctctccac ccttctcaga gtccagttt caaccagagt  
1381 tccaaccaat gggctccatc ctctggattc tggccaatga aatatctccc tggcagggtc  
1441 ctcttctttt ccagagctc ctcccaacc aggagctcta gttaatggag agtcccagc  
1501 acactcggac gcttgtgctt ttgtctccac gcaaacggat aaataaaagc attggtggcc  
1561 tt

FIG. 5 B/2

Length: 381 amino acids  
>AAA52025

```
1  MPFGNTHNKF  KLNYPKEEEY  PDLSKHNNHM  AKVLTLELYK  KLRDKEIPSG  FTVDDVIQTG
61  VDNPGHPFIM  TVGCVAGDEE  SYEVFKELFD  PIISDRHGGY  KPTDKHKTDL  NHENLKGDD
121 LDPNYVLSSP  VRTGRSIKGY  TLPFHCSRGE  RRAVEKLSVE  ALNSLTGEFK  GKYYPLKSM
181 EKEQQQLIDD  HFQFDKPVSP  LLLASGMARH  WPDAPGIWHN  DNKSFLVWVN  EEDHLRVISM
241 EKGGMKEVF  RRFCVGLQKI  EEIFKKAGHP  FMWNQHLGYV  LTCPSNLGTG  LRGGVHVKLA
301 HLSKHPKFEE  ILTRLRLQKR  GTGAVDTAAV  GSVFDVSNAD  RLGSESEVEQ  QLVVDGVKLM
361 VEMEKKLEKG  QSIDDMIPAQ  K
```

FIG. 5 C





FIG. 5 D



heart tissue sample

FIG. 5 E

Length: 403 nt  
>61166  
1 ACTTTGAGAA GTTACTTTCT AATTACGTCA TGAGAACACA ACTTGTAATT AGCAACACTT  
61 CTGTCAGTCT AGATCACTTC TTCTGCAGAG AGCTTTTCAA CCAAGTTGGC ATCAACCAGC  
121 ACAATAAAGT TTTCACCTGTT TTACCTGTTT CCTGTATATG GTGTAATCAG TGAAGAAGAAAT  
181 GGCATTTCAC ATCCTAAATA ATACGGTGAA ACACGTGCTA AAAATTACTT AGATTTAACA  
241 GAATTGCAAT TAGGTTTGA CAATGTATT ACTTCAAGAC AATGTATTTT ATCAGGAAAAA  
301 AATATCTTGA AAGAAAGATC TCTGAAATTA TTTTTCATT GATACGCCCTT TTCGTGACA  
361 AAATTTTGGG GTGAAATGGA TGATGTTTAC TGATTGATT AGT

FIG. 6 A

```

Length: 4828 nt
>61166contig
1  ttttccaagt ggaagttcaa ttgtcttttat ttttcttata cagattcaga gaagtaaaaa
61  ccagtaacca actccaggta aaatgggttg atctgacga tttggctgca tactttcggt
121 acgtataaca ttctaactt aaaatagaaa ttgttatatt acaaaaacgta gaagtaaaat
181 tttaaaaagt taaagtacta gcacatatat gtgttaggaa aatgggtctct gtcaattgcc
241 catthttcca attaaattaa cctacgattt cctthtttta acagcttatt tttttcataa
301 aagttgtact ttgagaagtt actttctaata tacgtcatga gaacacaaact tgtaatttagc
361 aacacttctg tcagtctaga tcacttcttc tgcagagagc ttttcaacca agttggcatc
421 aaccagcaca ataaagtttt cactgtttta cctgtttcct gtatatggtg taatcagtga
481 aagaaatggc atttcacatc ctaaatataa cggtgaaaca ctgtctaaaa attacttaga
541 tttaacagaa ttgcaattag gtthtgacaa tgtatttact tcaagacaat gtattttatc
601 aggaaaaaat atcttgaaag aaagatctct gaaattattt ttcatttgat acgccttttc
661 tgtgacaaaa ttttggggtg aaatgatgat gtthgctgat tgatttagta ctaaaaagac
721 tagtactaag aagactaaag acagttatct tataataaga aatatagtat aaatagcacc
781 ttatcaagaa ttctgcaggg gtthtaacac ttacaataat aggaataatgc cattaaaaag
841 ttgctctaac tttagatttc taactttagt gtthttaac aaaggccata tttgtggcc
901 ttaaaaaaaa aaaattatat ctggctttat ctattagtaa acacaaaagg tccatattht
961 attctgaaaa aatatthatt atattcattc ataaatgttc taactaattt aactaaaaaa
1021 atcttctagt atthtctgat gccacaagct tactagaaaa ttacttctaa aaatgggtaa
1081 tataaatcat caatgattha cctactthta aaaagagggg tatctgtttc tcttacattt

```

FIG 6 B/I

1141 aataaacctga aaatgagct ataaaaatat ttttaaaaaa tacagtaaca ctgctgagtt  
1201 ttgttaggtc ccttggtttt ttaatttttt atttatttat ttatttttag caagaatgta  
1261 caattctttt tgcaattttt tgctaacaaa agacaaaaag aaatagtgct cccttcaatt  
1321 tagtagcaat aaaatcatct atcttcatct ctctcagagg gcttagggag agtgaaaagg  
1381 attagaggaa cataaaccat gggtccttca ggtaaaataa gtcatttcat agtgatggag  
1441 gcaacagcag gctacgtctt tgtctgcctg tacgctcagc atcagcatgc cctgccccctc  
1501 ctccctccac ataagggtgca ccaaagcctt cttttcttct acaacagagt ctcccggtat  
1561 aaagggtggca acagaaaatt gttgctttgt ccttttgggg catctcttgc atctcctgct  
1621 tgtttttgctt ttgggggtact ataaacttgtc tctttccatt tcacaaactag tatcctttcc  
1681 acgatatctc ataaacttgc tacttcatct ctgacctttt cacttgacct ctttccaaca  
1741 tgatccaatg tactctgtac atgtatatct cggtagatca aaaggaaatct tatttaagat  
1801 cccaacaaa taagtcccc aatgcatttg atggaattg aaagtatcct ctcagagact caaattatta  
1861 aagtccttca aaaaagatta aatccataga ttataaataa tattagtcca aaatatataa  
1921 cagttgagga cttcattggc aatgcaggca gactgcatgc cagttgaaca tgatgctctc  
1981 tcagtcctta aaagctaatt aaaaatgggt ttggttacct aagagggtatt gaatacatat  
2041 ttcatgacct ttatatccaa ctgtagcaaa caggattagg ataataatct taggaatcaa  
2101 ttttactgaa ttcagaaaaca ttatatctca ccatacaccc tcaaagggca ttttttttta  
2161 catgtcagtc agagatctgc ttcatccttc agtttcatag atagaattat tttaaacact  
2221 tgaaatctag gaagcaaac tgacaaggct tcagaattta aaagcaaca gctcactgtg  
2281 tgtgggtgtgc tatcagggtg aaatctatgt tgtcctgatg ttttcagagt tatttcaaaa  
2341 gacaaaaata cagttgccac tgatttatca aaaacattg gctgcctttt gtcacagct

FIG. 6 B/2

```

2401 acaaaattac agtgctttat aaaataaaca tcaaggccgg gtgcggtggc tcacgccagt
2461 aatcccagca ctttggaagg ccgagatggg tggatcacct gagatcagga gtttgagaac
2521 agcctgacca ttatggtgaa accccgtctc tactcaaat acaaaaaatt agcgggacgt
2581 ggtggcaggc gcctgtaatc ccagctactc gggaggctaa ggcaggagaa tcgcttgaa
2641 ccagaaggca gaggttgagc tgagccgaga ttgtgccatt gcattccagc ctggacaaca
2701 agagcaaaac tcgatctcaa aaacaaacaa aacacatcaa gaaaaaaa aaaccatcag
2761 attctaagct gcaatttttt aaatccccag ttgtaatat tcaaaaactt ttgtttgaat
2821 aaatgctca taattagtac caactggtc tcttcacaa gatctgtagt gtaagaactg
2881 tgactaatgc tgctgctgct actttcacgc tggagggact aagtgtcta gatgctctgc
2941 agtaaaaata cgaagataa gctacaatag gactgtgtgc cttataaat acagactaat
3001 aagagccatc agagccagca tggattcaaa attacattgt attccataca gtagaatttt
3061 actatccata caatgatttt taaagctcaa gttaaatagt ttttaaagca ttgtggtacta
3121 ctgtcatcaa tacagttttt gaaactgtaa atcagggtcga attttgtgca catttcctgg
3181 accaagatgc ctcagaagt tccaattaac accataaag cttagccatg ggcagcaga gaagaaagag
3241 gtgtataaat tcaaatcaac accataaag cttagccatg ggcagcaga ggcagcaga
3301 aaaagtattc tgcataatca atcctgcaga cacaattctg tataatctgt cacaagaacg
3361 caggcttgca gaaaatgaaa atagaatatc tatttatgtt taacttaagt tactctcaat
3421 caaaaccagg caatgattaa actggcaaca taaaaggag ggagcacgag tcatggaggc
3481 gggaagtggc gcacctgcag acttgctctg ctccatcact ttttccaaga ggcccaggaa
3541 atgtaaggtc atggctacat ccaagtaca atggtagtga ttacagccag gttagaagg
3601 gctcactttt gttcagagca gactctacat cattgaagag ggggatcagg tcttcagatt

```

FIG. 6 B/3



```

3661 ccaaagtacc taagtcaacg tttgttcctg gaagacagtc aaggaaatca gggaaacggg
3721 tctgttgggg attgatgttc atgggtgttt gtcctgcgtt ttctcctgta tccatctcat
3781 ccacattgct cactgtcagt gctctgctcc ctggaatgat atggccctcc gtagcacccct aaccccaggc
3841 cactgtcagt tattattagt gatggatctc atgtcttgggg tcatcgtggg tgggttgaca gcagcctgaa
3901 ctggggcaag agtctcagct ttggcgcatc ttggtggga tccatggggg gctgtcgaca gagggcagct tcctgcctca
4021 tgagctcctc gtttctgctg ctgctgctgc ttggcgcatc cgaatccttt ctctctacat ctggattctc tgaagccgca
4081 gtttctgctg gtgggttctg agtgggtggg ctgctgctgc ttggtggga ttctgtggg gcgcatggg cagggccatg agccctgcgg
4141 gtgggttctg gtttctgctg agtgggtggg ttctgtggg ttctgtggg tcaggggtact gggggccatc tgctgctggg
4201 gttggtgatt gttggtgatt catcacgaga ttggtggg ttggtggg atactggcat ggacctctga ggcactggtg
4261 tggaactgac ggtcttgcca ggtggtgatt ggcaggggtgg aggttcatat aggtcagagg gattcagctc gccttcctag
4321 ggtcttgcca ggtggtgatt ggtggtgatt ggtggtgatt ggtggtgatt ggtggtgatt ggtggtgatt ggtggtgatt
4381 tgaaggctcat ctccagccc ggtggtgatt ggtggtgatt ggtggtgatt ggtggtgatt ggtggtgatt ggtggtgatt
4441 ggcggagggtg cgcgtgctgc ggcgtgctgc ggcgtgctgc ggcgtgctgc ggcgtgctgc ggcgtgctgc ggcgtgctgc
4501 gcagggacgc gcgggtggcc gcgggtggcc gcgggtggcc gcgggtggcc gcgggtggcc gcgggtggcc gcgggtggcc
4561 ccgggtggcc ccttaaagaa agactccggc gagggcttcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg
4621 ccttaaagaa agactccggc gagggcttcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg
4681 cagagttgaa gagggcttcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg
4741 gccagggcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg gagggcttcg
4801 aagctgagcc tgagcgcgcg gcggcgcggc gcggcgcggc gcggcgcggc gcggcgcggc gcggcgcggc gcggcgcggc

```

FIG. 6 B/4

Length: 398 amino acids  
>61166pep  
1 MNPASAPPPL PPPGQQVIHV TQDLDTDLEA LFNSVMNPKP SSWRKKILPE SFFKEPDSGS  
61 HSRQSSSTDSS GGHPPRLAG GAQHVRSHSS PASLQLGTGA GAAGSPAQQH AHLRQQSYDV  
121 TDELPLPPGW EMTFTATGQR YFLNHIEKIT TWQDPRKAMN QPLNHMNLHP AVSSTPVPQR  
181 SMAVSQPNLV MNHQHQQQMA PSTLSQQNHP TQNPAGLMS MPNALTQQQ QQQKLRLQRI  
241 QMERERIRMR QEELMRQEAA LCRQLPMEAE TLAPVQAAVN PPTMTDMS ITNNSSDPFL  
301 NGGPYHSREQ STDGGLGLGC YSVPTPEDF LSNVDEMDTG ENAGQTPMNI NPQQTRFPDF  
361 LDCLPGTNVD LGTLESEDLI PLFNDVESAL NKSEPFLLT

FIG. 6 C





FIG. 6 D



heart tissue sample

FIG. 6 E

Length: 168 nt  
>61244  
1 ACAGTTCCGG GAGGAACAAG ACCTTCCTCT GCTATGTGGT TGAAGCACAG GGCAAGGGGG  
61 GCCAAGTGCA GGCATCTCGG GGATACCTAG AGGATGAGCA TCGGGCTGCC CATGCAGAGG  
121 AAGCTTTCTT CAACACCATC CTGCCAGCCT TCGACCCAGC CCTGCGGT

FIG 7 A

Length: 1164 nt  
>AF161698

```

1  gaattccggc ggcctctctc ctctccctca gtgactcctg agccacagcc cctccatggc
61  ccagaaggaa gaggctgctg tggccactga ggctgcctcc cagaatgggg aggatctgga
121  gaacctggac gaccttgaga agctgaaaga gctgattgag ctgccgccct ttgagattgt
181  cacaggagaa cggctgcctg ccaacttctt taaattccag ttccggaatg tggagtacag
241  ttccgggagg aacaagacct tcctctgcta tgtggttgaa gcacagggca agggggggcca
301  agtgcaggca tctcgggat acctagagga tgagcatgcg gctgcccattg cagaggaagc
361  ttctctcaac acctcctgc cagccttcga ccagccctg ccgtacaatg tcacctggtg
421  tgtgtcctcc agcccctgtg cagcgtgtgc tgaccgcatt atcaaaaacc ttagcaagac
481  caagaaacctg cgtctgctca ttctggtggg tcgactcttc atgtgggagg agccggagat
541  ccaggctgct ctgaagaagc tgaaggaggc tggctgtaaa ctgcgcattca tgaagcccca
601  ggacttcgaa tatgtctggc agaattttgt ggagcaagaa gagggtgaat ccaaggcctt
661  tcagccctgg gaggacattc aggagaaactt cctatactac gaggagaagt tggcagacat
721  cctgaagtag ggcaactggg ctttgcctca cgtattcctg ctgccaccaa gagacagcaa
781  tgacatgtac agccatctgg gacatgcctg tcttcctaat accatttggg gctggacaa
841  atttgacacc aaccaatcat actggacaag gcccttagag gacttgaaat atacttctca
901  tgctgtagtt tatttaggct gtgactctct ctctaattgct gctctcggga aggacgaaag
961  tgacctgcaa ggagagaaat gcaaccatac atgggctcca gtcaactatg ggactgaagg
1021  tcctaattgc taccccaagg gggctgctta acacaaacag cctcagaccc gaggtttaga
1081  ttctctgaaat atgcatttta tgttaagtgt ggtatttttt taaaaaaga aaaacagcaa
1141  catataaaa agaagtgtg tggc

```

FIG. 7 B

Length: 224 amino acids  
>AAD45360  
1 MAQKEEAAVA TEAASQNGED LENLDDPEKL KELIELPPFE IVTGERLPAN FFKFQFRNVE  
61 YSSGRNKTFL CYVVEAQKG GQVQASRGYL EDEHAAAHAE EAFNTILPA FDPALRYNVT  
121 WYVSSSPCAA CADRIIKTLS KTKNLRLLIL VGRLFMWEEP EIQAALKKLK EAGCKLRIMK  
181 PQDFEYVWQN FVEQEEGESK AFQPWEDIQE NFLYYEEKLA DILK

FIG. 7 C




FIG. 7 D






FIG. 7 E

Length: 334 nt  
>65330  
1 ACTATCAATC TCCCTGGGAA CAAGCCATTA GCAATGATCC GGAGCTTTTA GAGGCTTTAT  
61 ATCCTAAACT TTTCAAGCCT GAAGGAAAGG CAGAACTGCC TGATTACAGG AGCTTTAACA  
121 GGGTTGCCAC ACCATTGGA GGTTTGAAA AAGCATCAAG AATGGTTAAA TTAAAGTTC  
181 CAGATTTTGA GCTACTATTG CTAACAGATC CCAGGTTTAT GTCCTTTGTC AATCCCCCTT  
241 CTGGCAGACG GTCCTTTAAT AGGACTCCCTA AGGGATGGAT ATCTGAGAAT ATTCCCTATAG  
301 TGATAACAAC CGAACCTACA GATGATACCA CTGT

FIG. 8 A

Length: 1590 nt  
>65330contig

```

1  gtcagagtag  ggaccatgct  gtcccaggtt  caaggataaa  aaccatcagg  cccaagtgcc
61  atccatagtc  catctccaga  gtcttcctcc  acaaaactggg  attcatcccc  gctgaaaaag
121 cacaatctaa  cagcaaggga  caagcaacag  ccatacatgaa  cataatacta  tgatgaagca
181 gagaaaaacag  caagcaacag  aaaaaggcca  gcatccccag  ggaaatgatg  ttgatggcat
241 ggacctgggc  aaaaaggcca  gcatccccag  agacatcatg  ttggaagaat  tatcccatct
301 cagtaaccgt  ggtgccaggc  tatttaagat  gcgtcaaaga  agatctgaca  aatacacatt
361 tgaaaaatttc  cagtatcaat  cttagagcaca  aataaatcac  agtatctgcta  tgcagaatgg
421 gaaagtggat  ggaagtaact  tggaaaggagg  ttgcgagcaa  gcccccttga  ctcttcccaa
481 caccgccagat  ccacgaagcc  ctccaaatcc  agacaacatt  gctccaggat  attctggacc
541 actgaaggaa  attcctcctg  aaaaattcaa  caccacagct  gtccctaagt  actatcaatc
601 tccctgggaa  caagccattt  gcaatgatcc  ggagctttta  gaggccttat  atcctaact
661 tttcaagcct  gaaggaaagg  cagaactgcc  tgattacagg  agctttaaca  gggttgccac
721 accatttgga  ggttttgaaa  aagcatcaag  aatggttaaa  tttaaaagttc  cagattttga
781 gctactattg  ctaacagatc  ccagggttat  gtcctttgtc  aatccccctt  ctggcagacg
841 gtcctttaat  aggactccta  agggatggat  atctgagaat  attcctatag  tgataaacac
901 cgaacctaca  gatgatacca  ctgtaccaga  atcagaagac  ctatgaaaag  aaagtgttat
961 gtgccacata  aaactctgaa  tataaaagtt  gctgttctac  tattttaact  actggcaaaag
1021 ccacttgcat  ttttcattag  tagcaacaat  agcaatttag  tgattttcct  tttctgacat
1081 tcaatttcaa  tctcagatca  aatactaata  aacaatttag  aatcttactt  taaaaaactt

```

FIG. 8 R/1

1141 ataactcact tgtcttcatt cataattttg tttcacctg gtttaaagaa tccagatat  
1201 ttactgcaaa agttcagatg gaaaagtaat tgacagyttc acctttgtct cattttatat  
1261 gattttattac agtgtaagtt tttcaagtgg aatctagaat caaaatcacag ggagagatat  
1321 gaagacctat tcagagtttc atctggggat gaaagctatg gaagatgatg tacaaatgtt  
1381 attgatggag aaaatggttg gtgtgtcctt tctggtgacc atgagaaaata tatgtcttga  
1441 tgaagtcttt tcattagtca ctcttagaat tctaaagtgc tttgcacttt tncaatatgt  
1501 ttctgaatcat taggtaattt attctgggnt gatattcttc aaaattcaat tcagttatta  
1561 tattcattta ggcattaagt caaggngact

FIG. 8 B/2

Length: 264 amino acids

>AAF63623

```
1  MLSHNTMMKQ  RKQQTAIMK  EVHGNDVDGM  DLGKKVSIPT  DIMLEELSHL  SNRGARLFKM
61  RQRRSDKYTF  ENFYQSRQAQ  INHSIAMQNG  KVDGSNLEGG  SQAPLTPPN  TPDPRSPPNP
121 DNIAPGYSGP  LKEIPPEKFN  TTAVPKYYQS  PWEQAISNDP  ELLEALYPKL  FKPEGKAELP
181 DYRSFNVRVAT  PFGGFEEKASR  MVKFKVPDFE  LLLLTDPFRM  SFVNPLSGRR  SFNRTPKGWI
241 SENIPIVITT  EPTDDTTVPE  SEDL
```

FIG. 8 C





FIG. 8 D



heart tissue sample

FIG. 8 E

Length: 290 nt  
>66214  
1 ACTCCTGAAG TGGAGGAGGG TGTTCCTCCC ACCTCGGATG AGGAGAAGAA GCCAATTCCA  
61 GGAGCGAAGA AACTTCCAGG ACCTGCAGTC AATCTATCGG AAATCCAGAA TATTAAAAGT  
121 GAACTAAAAT ATGTCCCCAA AGCTGAACAG TAGTAGGAAG AAAAAAGGAT TGATGTGAAG  
181 AAATAAAGAG GCAGAAAGATG GATTCAATAG CTCACTAAAA TTTTATATAT TTGTATGATG  
241 ATTGTGAACC TCCTGAATGC CTGAGACTCT AGCAGAAATG GCCTGTTTGT

FIG. 9 A



Length: 886 nt  
>66214cds

```

1  gtctctcaata cgggagagg cacagagcta ttccagccac atgaaaagca tcggaattga
61  gatcgagct  cagaggacac cgggcgccc ttccacctc caaggagctt tgtattcttg
121 catctggctg cctgggactt cccttaggca gtaaacaaat acataaagca gggataagac
181 tgcattgtata tgtcgaaaca gccagtttcc aatgttagag ccaccaggc aaatatcaat
241 attccaatgg gagcctttcg gccaggagca ggtcaacccc ccagaagaaa agaattgtact
301 cctgaagtgg aggaggggtgt tcctcccacc tcgcatgagg agaagaagcc aattccagga
361 gcgaagaaac ttccaggacc tgcagtcaat ctatcggaat tccagaatat taaaagtga
421 ctaaaatatg tccccaaagc tgaacagtag taggaagaaa aaaggattga tgtgaagaaa
481 taaagaggca gaagatggat tcaatagctc agactctagc tgtttgtaca ttatatctc
541 gtgaacctcc tgaatgcctg ttcttacttt atcttcattt ttggcacctc acagaacaaa
601 ttccctctag ttggctgtat ttcttacttt atcttcattt ttggcacctc acagaacaaa
661 ttagcccata aattcaacac ctggagggtg tggttttgag gaggatatg attttatgga
721 gaatgatatg gcaatgtgcc taacgatttt gatgaaaaagt tcccaagct acttcctaca
781 gtatttttgg caatatattg aatgcgtttt agttcttcac cttttaaatt atgtcactaa
841 acctttgtatg agttcaata aatatttgac taaatgtaaa atgtga

```

FIG. 9 B

Length: 88 amino acids  
 >66214pep  
 1 MYMSKQPVSN VRAIQANINI PMGAFRPGAG QPPRRKECTP EVEEGVPPTS DEEKKPIPGA  
 61 KKLPGPAVNL SEIQNIKSEL KYVPKAEQ

FIG. 9 C





FIG. 9 D



heart tissue sample

FIG. 9 E

Length: 152 nt, 59 nt and 234 nt  
>66268  
1 CTGATTATCA CAGCCCTCTT TTCTCCTGAA TTTTAAATGC AGAAGTTTGA ATGAAGCAAG  
61 GGAAGGCATG TAGGGACAGG AAAGGAAACA ATGGAAGGAA AGTGATTCTG TGAAAAGGAC  
121 AGTGAAGCCA GCTATTTTAC CCCCAGGCTG GA  
  
>52474  
1 TCCAGGGATT CCTTCCACGA CAGAAAACA TACAAGACTC CTTCAGCCAA CATGATGGT  
  
>S1MC01-1  
1 TCCANGGATT CCTTCCACGA CAGAANAACA TACAAGACTC CTTCAGCCAA CATGATGGTA  
61 CTGAAAGTAG AGGAACTGGT CACTGGAAAG AACAAATGGCA ATGGGAGGC AGGGAAATTC  
121 CTTCCCTGAGG ATTTCAGAGA TGGACAGTNT GAAGCTGCTG TTACTTTAGA GAAGCAGGAG  
181 GATCTGAAGA CACTTCTNGC CCACCCTGTG ACCCTGGGG AGCAACNGTG GAAA

FIG. 10 A

Length: 1901 nt  
 >X83703

```

1  aaaaaacagc  agggttagct  tgtccctccc  ctccctcttc  agcttcccag  aactgattc
61  tggaatgaaa  attcacctgc  ctctgagttg  gctcctaata  ggggtgggag  tgttacttcg
121  gttcccaggt  tggaagatta  tctcacccgg  cccagctat  ataagctgac  cgggtgggag
181  gggcccagca  gggccaactc  cagggatccc  ttcacgaca  gaaaaacata  caagactcct
241  tcagccaaca  tgatggtact  gaaagtagag  gaactggtca  ctggaaagaa  gaatggcaat
301  ggggaggcag  ggaattcct  tctgaggat  ttcagagatg  gagagtatga  agctgctgtt
361  actttagaga  agcaggagga  tctgaagaca  ctctagccc  accctgtgac  cctgggggag
421  caacagtgga  aaagcgagaa  acaacgagag  gcagagctcc  caaagaaaaa  actagaaaca
481  agatccaagc  ttgaaaattt  agaagacctt  gaaataatca  ttcaactgaa  gaaaaggaaa
541  aaatacagga  aaactaaagt  tccagttgta  aaggaaaccag  aacctgaaat  cattacggaa
601  cctgtggatg  tgcctacgtt  tctgaaggct  gctctggaga  ataaactgcc  agtagtagaa
661  aaattcttgt  cagacaaaga  caatccagat  gtttgtgatg  agtataaacg  gacagctctt
721  catagagcat  gcttggaagg  acatttggca  atttgtgaga  agttaatgga  agctggagcc
781  cagatcgaat  tccgtgatat  gcttgaatcc  acagccatcc  actgggcaag  ccgtggagga
841  aacctggatg  ttttaaaatt  gttgctgaat  aaaggagcaa  aaattagcgc  ccgagataag
901  ttgctcagca  cagcgctgca  tgtggcgggtg  aggactggcc  actatgagtg  cgcggagcat
961  cttatcgcct  gtgaggcaga  cctcaacgcc  aaagacagag  aaggagatag  cccgttgcat
1021  gatgcggtga  gactgaaccg  ctataagatg  atccgactcc  tgattatgta  tggcgcggat
1081  ctcaacatca  agaactgtgc  tgggaagacg  ccgatggatc  tggtgctaca  ctggcagaat

```

FIG. 10 B/1

```

1141 ggaaccaaag caatattcga cagcctcaga gagaactcct acaagacctc tcgcatagct
1201 acattctgag gcaaacgaca gactctttaat cagtaaatgt tcactggcat tttgaaggca
1261 tggcccagga gaagagacac tagccataaa atctagtttc tatttatcaa cgtgttgtga
1321 agatgtacct aatgaagtgt tgagaaagca cagggttata ggtgtttaaa tttccttttag
1381 tgaactctt atttattttt atgtattcct gtttatttat ttactgccac gctactgata
1441 ttcagaccct catgattcat catctggtga gcagagcttc atttgtatat aacactttca
1501 gaggcttccc accataggt agttcttaaa ccaggtgaaa gagcaaatgt caagtgccta
1561 ctatgtgtc attcgtcat gtaagagttt ttaagagagg gctgattatc acagccctct
1621 tttctcctga atttttaatg cagaagtttg aatgaagcaa gggaaggcat gtagggacag
1681 gaaaggaaac aatggaaagg aagtgattct gtgaaaaagg cagtgaagcc agctatttta
1741 ccccaggct ggattttttt tttttttttt tttttttttt tttttaccga gtacacagag
1801 tacccaagtg aagagaacgt catgagtgtg agtgcaaatc agtggaagga gcggcaaac
1861 gggacatgca gaattgaatt tgctcaaaaa aaaaaaaaaa a

```


FIG 10 B/2

Length: 319 amino acids

>CAA58676

```
1  MMVLKVEELV  TGKKNNGEA  GEFLPEDFRD  GEYEAAVTLE  KQEDLKTLLA  HPVTLGEQQW
61  KSEKQREAEL  PKKLEQRSK  LENLEDLEII  IQLKKRKKYR  KTKVPVVKEP  EPEIITEPVD
121  VPTFLKAAL  E  NKLPVVEKFL  SDKNNPDVCD  EYKRTALHRA  CLEGHLAIVE  KLMEAGAQIE
181  FRDMLESTAI  HWASRGNLD  VLKLLLNKGA  KISARDKLLS  TALHVAVRTG  HYECAEHLIA
241  CEADLNAKDR  EGDTPLHDAV  RLNRYKMIRL  LIMYGADLNI  KNCAGKTPMD  LVLHWQNGTK
301  AIFDSLRENS  YKTSRIATF
```

FIG. 10 C



66268 (152 nt)

52474 (59 nt)

S1MC01-1 (234 nt)

X83703 (1901 nt)

or 66268cds

CAA58676 (319 aa)

or 66268pep

FIG. 10 D






FIG. 10E




FIG. 10 F

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number  
**WO 01/092567 A3**

(51) International Patent Classification: C12Q 1/68.  
A01K 67/027, A61K 49/00, A61P 9/00, C07K 16/18.  
G01N 33/53

Birgit [DE/DE]: Heiterwangerstrasse 10, 81373 Munich (DE). BECK, Joachim [DE/DE]; Henerichstrasse 115, 81477 Munich (DE). HENKEL, Thomas [DE/DE]; Freienfelsstrasse 20a, 81249 Munich (DE).

(21) International Application Number: PCT/EP01/06165

(74) Agents: LEIDESCHER, Thomas et al.; Zimmermann & Partner, Postfach 33 09 20, 80069 Munich (DE).

(22) International Filing Date: 30 May 2001 (30.05.2001)

(25) Filing Language: English

(81) Designated States (national): AU, CA, JP, US.

(26) Publication Language: English

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(30) Priority Data:  
60/207,400 30 May 2000 (30.05.2000) US

Published:  
--- with international search report

(71) Applicant (for all designated States except US): MEDI-  
GENE AG [DE/DE]; Lochhamer Strasse 11, 82152  
Planegg/Martinsried (DE).

(88) Date of publication of the international search report:  
13 February 2003

(72) Inventors: and

(75) Inventors/Applicants (for US only): BUNK, Daniela  
[DE/DE]; Hofmark 8, 82393 Iffeldorf (DE). REUNER,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/092567 A3

(54) Title: NOVEL TARGET GENES FOR DISEASES OF THE HEART

(57) Abstract: The present invention relates to a variety of genes abnormally expressed in heart tissue as well as to fragments of such genes. Assessment of the expression level of these genes may be used for testing the predisposition of mammals and preferably humans for a heart disease or for an acute state of such a disease. Preferred diseases in accordance with the invention are congestive heart failure, dilative cardiomyopathy, hypertrophic cardiomyopathy and ischemic cardiomyopathy. The present invention further relates to methods of identifying compounds capable of normalizing the expression level of the aforementioned genes and of further genes affected by the abnormal expression. The identified compounds may be used for formulating compositions, preferably pharmaceutical compositions for preventing or treating diseases. They may also be used as lead compounds for the development of medicaments having an improved efficiency, a longer half-life, a decreased toxicity etc. and to be employed in the treatment of heart diseases. Included in the invention are also somatic gene therapy methods comprising the introduction of at least one functional copy of any of the above-mentioned genes into a suitable cell. Finally, the invention relates to non-human transgenic animals comprising at least one of the aforementioned genes in their germ line. The transgenic animals of the invention may be used for the development of medicaments for the treatment of heart diseases.

## INTERNATIONAL SEARCH REPORT

Inter of Application No.

PCT/EP 01/06165

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C1201/68 A01K67/027 A61K49/00 A61P9/00 C07K16/18  
G01N33/53

According to International Patent Classification (IPC) or to both national classification and IPC.

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols):

IPC 7 C12Q G01N A61K A61P C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! EBI; 3 May 2000 (2000-05-03) WALKER M ET AL: "Cardiomyopathy-associated genes identified by genome-scale expression analysis" Database accession no. AW755252 XP002203686 abstract</p> <p>---</p> <p>-/--</p>	1-41,47

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search:

18 October 2002

Date of mailing of the international search report

28. 10. 2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Osborne, H

## INTERNATIONAL SEARCH REPORT

Intern I Application No

PCT/EP 01/06165

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>AIHARA Y ET AL: "CARDIAC ANKYRIN REPEAT PROTEIN CARP IN A NOVEL MARKER OF MYOCARDIAL STRESS: REGULATION OF TRANSCRIPTION THROUGH M-CAT ELEMENT BY STRESS-RESPONSIVE MAP KINASE PATHWAYS" CIRCULATION, AMERICAN HEART ASSOCIATION, DALLAS, TX, US, vol. 110, no. 18, SUPPL, 2 November 1999 (1999-11-02), pages I-627-I-628, XP001056080 ISSN: 0009-7322</p>	1-41,47
Y	<p>WO 99 24571 A (CURAGEN CORP ;LOWE DAVID G (US); SHIMKETS RICHARD A (US)) 20 May 1999 (1999-05-20) the whole document</p>	1-41,47
Y	<p>WO 99 49062 A (FAN WUFANG ;GENE LOGIC INC (US); PRASHAR YATINDRA (US)) 30 September 1999 (1999-09-30) claims 17-22</p>	1-41,47
Y	<p>DE 197 25 186 A (MEDIGENE AG) 24 December 1998 (1998-12-24) page 7, line 55 -page 8; claims 6-15</p>	1-41,47
Y	<p>WO 99 50410 A (INST MEDICAL W &amp; E HALL ;VICTOR CHANG CARDIAC RESEARCH (AU); HARVE) 7 October 1999 (1999-10-07) the whole document</p>	1-41,47
Y	<p>LOWES B D ET AL: "CHANGES IN GENE EXPRESSION IN THE INTACT HUMAN HEART. DOWNREGULATION OF ALPHA-MYOSIN HEAVY CHAIN IN HYPERTROPHIED, FAILING VENTRICULAR MYOCARDIUM" JOURNAL OF CLINICAL INVESTIGATION, NEW YORK, NY, US, vol. 100, no. 9, 1 November 1997 (1997-11-01), pages 2315-2321, XP002910970 ISSN: 0021-9738 the whole document</p>	1-41,47
Y	<p>KELLER ET AL: "Differential expression of alpha- and beta-enolase genes during rat heart development and hypertrophy" AMERICAN JOURNAL OF PHYSIOLOGY: HEART AND CIRCULATORY PHYSIOLOGY, THE AMERICAN PHYSIOLOGICAL SOCIETY, XX, vol. 38, no. 6, 1 December 1995 (1995-12-01), pages H1843-H1851, XP002096499 ISSN: 0363-6135 the whole document</p>	1-41,47

-/-

## INTERNATIONAL SEARCH REPORT

Intern Application No

PCT/EP 01/06165

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BOHELER ET AL: "Gene expression in cardiac hypertrophy" TRENDS IN CARDIOVASCULAR, ELSEVIER SCIENCE, NEW YORK, NY, US, vol. 2, no. 5, 1 September 1992 (1992-09-01), pages 176-182, XP002096139 ISSN: 1050-1738 -----	1-41,47
Y	WO 91 01498 A (VIOCLONE BIOLOG INC) 7 February 1991 (1991-02-07) the whole document -----	40,41,47
A	WO 99 19471 A (MOLKENTIN JEFFREY D ;GRANT STEPHEN R (US); OLSON ERIC N (US); UNIV) 22 April 1999 (1999-04-22) -----	
A	WO 98 13476 A (LEINWAND LESLIE A ;VIKSTROM KAREN L (US)) 2 April 1998 (1998-04-02) -----	

# INTERNATIONAL SEARCH REPORT

onal application No.  
PCT/EP 01/06165

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 35-38, and 41 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

As a result of the prior review under R. 40.2(e) PCT,  
all additional fees are to be refunded.

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:  
  
1-41,47 with respect to SEQ ID Nos 1 and 10
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☒ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-41,47 all partially

Methods for screening, diagnosing of individuals for the presence of a heart disorder based on the differential expression of gene/gene product markers; methods for identifying compounds and uses thereof in the treatment of heart disorders, wherein the first gene marker is SEQ ID NO 1 i.e. (MYOM2).

2. Claims: 1-39,47 all partially

Methods for screening, diagnosing of individuals for the presence of a heart disorder based on the differential expression of gene/gene product markers; methods for identifying compounds and uses thereof in the treatment of heart disorders, wherein the first gene marker is SEQ ID NO 2.

3. Claims: 1-39,47 all partially

Methods for screening, diagnosing of individuals for the presence of a heart disorder based on the differential expression of gene/gene product markers; methods for identifying compounds and uses thereof in the treatment of heart disorders, wherein the first gene marker is SEQ ID NO 3.

4. Claims: 1-47 all partially

Subjects 4-19 IDEM, wherein each of the subjects relates to a distinct SEQ ID NO, i.e. SEQ ID NO 4 (and to the muscle isoform of creatine kinase); SEQ ID NO 5 (and YAP65); SEQ ID NO 6 (and APOBEC-2); SEQ ID NO 7; SEQ ID NO 8 (and SMPX/Cs1); SEQ ID NO 9 ( and C-193/CARP); SEQ ID NO 10; SEQ ID NO 11 etc ... SEQ ID NO 19.



## INTERNATIONAL SEARCH REPORT

Inter. Application No  
PCT/EP 01/06165

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9924571	A	20-05-1999	AU 1395399 A	31-05-1999
			CA 2309360 A1	20-05-1999
			EP 1030916 A2	30-08-2000
			JP 2002508923 T	26-03-2002
			WO 9924571 A2	20-05-1999
WO 9949062	A	30-09-1999	AU 3208999 A	18-10-1999
			CA 2323574 A1	30-09-1999
			EP 1073756 A1	07-02-2001
			WO 9949062 A1	30-09-1999
DE 19725186	A	24-12-1998	DE 19725186 A1	24-12-1998
			AU 746190 B2	18-04-2002
			AU 8438598 A	30-12-1998
			WO 9856907 A1	17-12-1998
			EP 1000145 A1	17-05-2000
			JP 2002509433 T	26-03-2002
WO 9950410	A	07-10-1999	AU 3127799 A	18-10-1999
			WO 9950410 A1	07-10-1999
			EP 1064374 A1	03-01-2001
WO 9101498	A	07-02-1991	AU 5960890 A	22-02-1991
			WO 9101498 A1	07-02-1991
WO 9919471	A	22-04-1999	AU 1099299 A	03-05-1999
			AU 748334 B2	30-05-2002
			AU 9805898 A	03-05-1999
			CA 2306448 A1	22-04-1999
			CA 2315244 A1	22-04-1999
			EP 1023441 A1	02-08-2000
			EP 1025216 A1	09-08-2000
			JP 2001520170 T	30-10-2001
			JP 2001520010 T	30-10-2001
			WO 9919471 A1	22-04-1999
			WO 9919473 A1	22-04-1999
			US 6201165 B1	13-03-2001
			US 2002056144 A1	09-05-2002
WO 9813476	A	02-04-1998	AU 4503497 A	17-04-1998
			EP 1007628 A1	14-06-2000
			WO 9813476 A1	02-04-1998
			US 6353151 B1	05-03-2002